

International Commission for



the Northwest Atlantic Fisheries

Serial No. 4016
(D.c.11)

ICNAF Res.Doc. 76/X/130

SPECIAL MEETING OF PANEL A (SEALS) - OCTOBER 1976

A stochastic model for the management of the Northwestern
Atlantic harp seal *Pagophilus groenlandicus* population

by

T. Berjaminsen
Institute of Marine Research
Directorate of Fisheries
N-5011 Bergen-Nordnes, Norway

and

P.F. Lett
Marine Fish Division
Bedford Institute of Oceanography
Dartmouth, Nova Scotia, Canada

Advice from the scientific advisers under the auspices to ICNAF to the international commissioners for 1977 was that the TAC for harp seals should not exceed 170,000. This advice, in part, was based on the scientific arguments presented in this paper. A stochastic model is developed which takes into account the variations in natural mortality and the landmen, high Arctic and Greenland catches. The Canadian - Norwegian large vessel hunt is controlled under quota regulations. The model is non-linear a result of changes in fertility and fecundity rates in response to shifts in population size. The MSY 1+ population size is determined to be 1.6 million seals, or a breeding stock size of 375,000 seals. The MSY is approximately 240,000 seals assuming the hunt continues its present pattern. The 240,000 can further be split into 200,000 pups and 40,000 1+ seals. Present stock size is \approx 1.2 million and a TAC of 170,000 seals will allow the population size to reach to MSY level in 10 to 15 years. A number of other management strategies are considered, in addition to prospects for future research.

Introduction

The exploitation of the northwestern Atlantic harp seal has been well documented, especially in recent years. Nearly 10 million harp seals are known to have been harvested from this population between 1895 and 1946 (Fisher 1955) and another 5.3 million were taken between 1947 and 1964 (Department of Fisheries of Canada 1968; Øritsland 1967). From 1965 to 1974, another 2.3 million seals were slaughtered in the northwestern Atlantic. This gives an average minimum annual kill over the period of 220,000 seals; a figure which has sustained the fishery for 80 years. Recent regulations have set a quota for seals at 150,000 animals between 1972 and 1975, (ICNAF 1972) with a low quota restriction in 1976 only permitting the take of 127,000 seals. These quotas do not include the high Arctic or Greenland catches, which can be considered to be approximately 10,000 animals. However, despite the low quota for 1976, *174,000 seals were killed.

Conflicting views as to the present status of the northwestern Atlantic harp seal have recently generated much controversy. These divergent opinions resulted from the interpretation of incomplete data sets, and poor communication among participating scientists. However, there are methods such as ultra-violet aerial sensing, which it is thought can be used independent of supportive biological information, and do not rely on current scientific opinion. Unfortunately, to date this method produces a "best estimate" of pup production in 1975 which is approximately 15,000 animals less than the catch (Lavigne et al. 1975).

*This number is subject to slight alterations.

Due to large discrepancies in estimation of production and natural mortality, it has been necessary to revise the basic data to incorporate the effects of the diverse fisheries, in addition to use advanced statistical and computer techniques to assess the data; thus producing a reliable assessment of the northwestern Atlantic harp seal stock.

Materials and methods

AMALGAMATION OF DATA

The assessment of an animal population requires the estimation of certain vital rates. The estimation of these rates, to a large extent, consists of the analysis of age frequencies which either represent the population or the catch structure. In some populations the catch and population age structures are assumed synonymous, however, this assumption can lead to grave errors in the instance of the northwestern Atlantic harp seal population (Benjaminsen and Øritsland MS 1975).

Initially, the problem consists of producing an age frequency of the total annual catch for one year old and older seals (1+), which amalgamates the catch frequencies from the individual fisheries in their proper proportions. In this fishery the age distribution for the different fisheries is indeed diverse.

Shot samples from Notre Dame Bay, Newfoundland, consist primarily of bedlamers. The La Tabatière, Que., and Labrador net fisheries

usually result in a sample with a preponderance of mature animals, while the St. Anthony, Nfld., shot samples seem to be more representative for the population structure. It is assumed that on average each of these fisheries tend to be roughly equivalent to each other in overall catch and can be summed to represent the landmen catch without producing serious errors. The large ships catch from the moulting lairs, however, is quite different in structure from the overall landmen's and must be treated separately. One of the primary problems is the fraction of one year old seals in the sample. This age group is usually segregated from the remaining age groups and is only randomly encountered when sampling the moulting lair (Benjaminsen and Øritsland MS 1975). Furthermore, the closing date also effects the large vessel sample since the earlier the hunt in the moulting lairs is terminated, the less females are represented in the catch (Sergeant 1965, Øritsland 1971). Thus the mature composite age frequencies vary depending on the length of the hunt, and consist mainly of males (Fig. 1). Samples of the Greenland and the high Arctic hunt, representing 8% of the total catch on average, are excluded from this analysis from 1952 to 1975, since no consistent sampling and catch records are available.

Between 1952 and 1960 jaws were collected on a regular basis from the landmen's catch and this sector of the hunt can be considered to be well represented (Sergeant and Fisher 1960, personal communication) and are presented in Table 1. During the years 1952-54 and 1957-58 samples were also collected from the large vessels. However, the 1957-58 samples

are sparse and it is unlikely that they accurately represent the catch (Table 1). For this reason the years 1955 to 1960 were replaced by an average catch frequency for large vessels. The attendant errors are possibly serious since during this period the large vessel hunt on 1+ animals represented between 87.5-95.5% of the total catch (Table 2).

In 1961 sampling began to steadily improve with both the large vessel catch from the moulting lairs and the landsmen catch being well represented (Table 1). Annual catch at age samples from 1961 on came from a number of sources (Sergeant 1971, 1972, personal communication; Øritsland 1971; Benjaminsen and Øritsland MS 1975). Unfortunately, there was not a good sample taken for either the landsmen or the large vessel catch in 1972, but the available data were included in the ensuing analysis. In 1973 the landsmen sample was again lacking and may not sufficiently represent the fishery. In both 1972 and 1973 there was no basis for using an average sample to represent either fishery, especially since the landsmen and large vessel catches are nearly equivalent to one another. In general, the samples from large vessels seemed to improve from 1967 on, excluding 1972.

The samples from the landsmen hunt and large vessel hunt were first summed for each year, then reduced to their respective percentage compositions. These compositions were then weighted in accordance with the numbers of animals killed in the landsmen and large vessel hunt (summed across ages) and subsequently divided by the sum of weighting factors. In this way then, the age composition data gave

the best possible representation of the catch distribution as the hunt has shifted from large vessels to landsmen (Fig. 2). Using this catch distribution the total catch was broken out into catch at age (Table 3).

Biological Basis of the Model

CALCULATION OF NATURAL MORTALITY

The instantaneous rate of natural mortality is possibly the most elusive parameter in population dynamics to estimate. Furthermore, for the harp seal population it is the most important due to the rather low exploitation rate experienced by 1+ animals.

The only representative sample of the population age distribution comes from males of age two and older in the moulting lairs (Benjaminsen and Øritsland MS 1975). Furthermore, it is assumed that there is no difference between natural mortalities rates of male and female harp seals. This is a valid assumption since male and female harp seals seem to experience similar growth rates and achieve equivalent maximum weights (Sergeant 1973a). The metabolic rate and body size of seals is well correlated (Lavigne et al. 1976); and since the mortality and metabolic rates of animals are related (Simms et al. 1959) it is unlikely that male or female harp seals experience divergent natural mortalities.

Male age samples of moulting seals were taken from Benjaminsen and Øritsland (MS 1975) for 1969, 1970, 1971, 1973 and 1974. In 1968, 1970 and 1973 there were also combined (male and female) samples which were corrected, (Fig. 1), for the fraction of males. These corrected

samples were added to the appropriate sexed male sample to produce a total male sample. The age frequencies were then reduced to percentage age compositions so the total instantaneous mortality rates (Z) could be calculated. Total instantaneous mortality rates, weighted for varying cohort abundance, were calculated by using the equation:

$$(1) Z_{(t,t+1)} = \log_e \left[\frac{\sum_{i=2}^{i=24} p_t}{\sum_{i=3}^{i=25} p_{t+1}} \right]$$

where P is the percentage in year (t) for age (i). One year old animals were not included because of the erratic nature in which they occurred in the sample.

Since there was not a representative male sample in 1972 the total mortality was calculated from 1971 to 1973 and the result value divided by two.

The weighted annual instantaneous mortality rates were as follows:

| Year | Total mortality (z) | |
|-----------|---------------------|---------------------|
| 1968-1969 | 0.0667 | } $\bar{z} = 0.160$ |
| 1969-1970 | 0.2247 | |
| 1970-1971 | 0.1882 | } $\bar{z} = 0.137$ |
| 1971-1973 | 0.0892 (Avg.) | |
| 1973-1974 | 0.1836 | |

The following two simultaneous equations could then be constructed and solved.

$$\bar{F}_{68-71} + M = 0.160$$

$$\bar{F}_{71-74} + M = 0.137$$

where F is the average instantaneous rate of hunting mortality.

Following the implementation of the quota regulations in 1971, hunting effort exerted on 1+ seals was essentially halved such that

$$F_{68-71} = 2F_{71-74}$$

Hunting effort was calculated by multiplying the number of men by the length of time they hunted 1+ seals and the mean horse power of the vessels. This value was then divided by the mean tonnage of the vessels, the number of days was calculated according to the closing dates of the hunt and the other values are taken from ICNAF statistical yearbooks. Ice condition is not considered in the calculation because of the lack of data, however, its inclusion would presumably greatly improve the calculation of effective effort. The hunting effort, which varies little from year to year was 120,000 men, day, hp, ton⁻¹ between 1961 and 1970, and 56,000 between 1972 and 1975. In both cases the standard deviation was only 8900 man, hp, ton⁻¹. Thus the change in effort is 2.07 and the assumption of a halving of effort between the two periods seems quite valid. The two equations then give the following values:

$$M = 0.114 \text{ and } \bar{F}_{68-71} = 0.046 \text{ and } \bar{F}_{71-74} = 0.023.$$

By subtracting the appropriate F value from the z values a standard deviation of 0.0677 and standard error of 0.0302 were calculated. From this limited sample set the variance was indeed great and it is assumed that the standard error more closely reflects the real biological deviation in natural mortality. Other calculations of mortality are somewhat lower, near 8% per year (Ricker 1971, Ulltang 1971).

Lavigne et al. (1976) present an age specific natural mortality schedule which declines from 0.2 for 0-group seals to 0.095 for 5 to 6 year olds. Following this, mortality rises to 0.109 for adults and remains constant. This schedule, although assumed, makes good biological sense since mortality and growth are usually coupled (Simms et al. 1959, Bourlière 1959). For harp seals the growth rate begins to become constant at the onset of maturity. According to an analysis of Lavigne's et al. (1976) data the instantaneous growth rate of mature seals is 0.12, not far from our estimated mortality of 0.114. However, at some point the mortality rate must exceed growth rate or the biomass of the cohort of seals would continue to increase forever. Senescent death in harp seals may begin at approximately age 18, if one speculates as to the reason about inflection in the survivorship curve at this time. Thus the critical age for a cohort of seals probably is near 18 years of age. In our analysis we could find no evidence of an age dependent natural mortality rate, which is not surprising considering the crudity of catch

data and the delicate changes in the parameter as suggested by Lavigne et al. (1976). However, the analysis was on 2+ seal and a different mortality rate may be experienced by the younger animals.

SEQUENTIAL POPULATION ANALYSIS

Sequential population methods (Fry 1949; Murphy 1964; Jones 1964; Gulland 1965; Pope 1972; Doubleday 1975) were developed to estimate fish population sizes and fishing mortalities from catch-at-age data when effort data are not available. Although there are no examples of this method being used to assess mammalian populations, no assumptions are violated by its use. For this reason it may be instructional to outline the method.

The sequential population analysis developed by Pope (1972) called cohort analysis was utilized in this study since it assumes that natural and hunting mortality occur somewhat seasonally, which is true to a great extent in the seal fishery. The method is based on the formula:

$$(2) \quad N_i = C_i \text{EXP } M/2 + N_{i+1} \text{EXP } M$$

where N_i is the population of a year-class at the i^{th} birthday, C_i is the catch of a cohort at age i , and M is the instantaneous coefficient of natural mortality. This formula is applied sequentially, the population size is each year depending on the population the year after.

However, some starting values are required. Thus, by expanding equation (2):

$$(3) N_i = (C_i \text{EXP}(M/2) + C_{i+1} \text{EXP}(3M/2) + C_{i+2} \text{EXP}(5M/2) + \dots + N_t \text{EXP}((t-i)M)$$

and assuming that hunting does not completely extirpate a particular cohort, the last term for the final year's population is:

$$(4) N_i = \frac{C_i (F_{i+1} + M)}{F_i (1 - \text{EXP}(-F_{i+1} - M))}$$

Equation 4 then, is used to calculate the population size in the initial year. Thus one of the primary problems is to estimate starting hunting mortality values.

The average hunting mortality calculated for 1972-74 was 0.023 and it was assumed that this value also applied in 1975. The analysis was not started using the 1976 data since all samples are not yet completely analysed. When possible considerable care should be given in estimating the initial hunting mortality since hunting mortalities are low for harp seals and the analysis is therefore more sensitive to initial F for a longer period. However, when the data series is long, poor starting values of F are considerably improved as the analysis continues. After running the analysis and averaging hunting mortalities

for ages 10 to 20 it was determined that on average a hunting mortality of 0.02 was adequate for animals of age 25. The age specific hunting mortalities calculated by the formula

$$(5) F = \text{Log}_e (N_i / N_{i+1}) - M$$

are presented in Table 4.

The resultant population estimates (Table 5) indicate that there was a quite dramatic decline, such that in 1968 the herd was only 45% as large as its size in 1952. Pup productions estimated by cohort analysis are somewhat erratic, indeed more than would be expected (Fig. 3) for a mammalian population. This inconsistency can perhaps be attributed to a natural mortality rate among pups which responds in some way to the exploitation rate. Furthermore interspecific competition may be a factor in determining resultant natural mortality rates.

DENSITY DEPENDENT PREGNANCY AND WHELPING AGE

It has been proposed by Sergeant (1966, 1973b) that the mean age of whelping for harp seals is a density dependent function relying on population size. This phenomenon is well noted in some other marine mammals (Gambell 1973). Indeed, some sort of mechanism is necessary to equilibrate the population with the carrying capacity of its environment (McLaren 1967). Sergeant (1976) has presented further evidence that the mean age of whelping has shifted to an extremely low level of

4.8 years in 1976, a figure well below the 6.5 years calculated in 1953. However, the reason for this most recent shift from 1968-1976 is not clear since the population has stabilized and seems to be slightly increasing. Furthermore, it will not become lucid until some information on the growth rate of individuals within cohorts becomes available. Sergeant (personal communication) has suggested that the maturity schedule of seals is determined by their growth rate as juveniles since growth approaches an asymptote at about age 5 (Sergeant 1973a). Lavigne et al.'s (1976) data would also suggest an age of 5. In both examples, the data are extremely variable. The harp seal population did reach it's minimum level between 1966 and 1973 of 1.07 million 1+ seals (Table 5). Assuming that the growth rate of juvenile seals is stock dependent, this would suggest that Sergeant's hypothesis is possibly correct. Until more conclusive evidence concerning the reproductive biology of harp seals is presented, we assume that the maturity schedule of harp seals is dependent on the coincident population size of 1+ animals. There is little evidence to suggest that the Front and Gulf herds experience a different maturity schedule, although this point may be argued by Sergeant (1973b), thus the data utilized from these two areas was not weighted in relation to population size.

The ogive were plotted on "probit" paper and lines were fitted by eye, giving more weight to points closer to the 50% maturity level. Values were then interpolated for each age from these lines, under the assumption that they represented the best fit of the data. The

interpolated values were used to determine the following equation (Fig. 4):

$$(6) \text{ ARC sine } {}_i E_t = 15.522A - 2.245 \times 10^{-5} N_t - 16.017$$

where ${}_i E_t$ is the fraction of the population whelping, assuming 100% pregnancy for a particular age i , N_t is the population number of $1+$ seals in the year t , and A is the age in years.

The shift in maturity in response to population size is assumed to be linear since the best fit of the mean age of whelping over the data series was linear, although again the data are not conclusive. Capstick and Ronald (1976) fit an exponential relationship to two data points, where there seems to be some confusion concerning the independent and dependent variables (see Fig. 4 Capstick and Ronald 1976). In fact, the curve has doubtful biological meaning since according to their hypothesis the mean age of maturity approaches zero at population sizes less than 1 million, a result which they previously speculate as being impossible. In addition, maturity reaches an asymptote at larger herd sizes, such that the population can increase ad infinitum. Surely the opposite effect is expected if it is indeed non-linear, since the biological basis for a shifting maturity ogive would be to constrain the population within the environmental carrying capacity.

Density dependent, age specific maturity would be enough to limit the population, but this is only one of a multitude of factors.

Others responding to population size could be natural mortality and fertility rate. Although we have no evidence for density dependent mortality, there does seem to be some data supporting a varying fertility rate (Fig. 5). Fertility rates were determined from a number of sources (Fisher 1952; Sergeant 1966, 1969, 1970, 1976; Øritsland 1971). The response seems to be a linearly decreasing function of population size, however, the data are variable and the structure of the relationship is not clear. The data were plotted against the 2+ population size since the younger animals remain segregated from the herd and would not compete. A fluctuating fertility rate is a well noted phenomenon and has been observed in at least three populations of whales (Gambell 1973). In addition, the unexploited population of Antarctic crabeater seals (*Lobodon carcinophagus*), which like harp seals also enjoy an unlimited ice substrate on which to whelp, have a low pregnancy rate of 0.76 (Øritsland 1970). Markgren (1969) found that the ovulation rate in moose (*Alces alces*) was related to a number of factors such as age, body size, nutrition, climate and population density.

EFFECTS OF CHANGING SEX RATIO ON POPULATION PROJECTION

The pup production estimates resulting from sequential population analysis are quite erratic (Fig. 3), possibly as a result of unaccountable fluctuations in natural mortality. In addition, the preponderance of males in the kill from the moulting seals would result in a sex ratio favouring females. This could increase the

estimates of pup production by 15,000 or 20,000 animals.

It is unlikely that there are serious errors in the catch distributions for seals 1 to 6 (Table 3), since only 8% of the variation of this data is unexplained when compared with the actual catch of bedlamers (see Table 2 Øritsland and Benjaminsen MS 1975). This comparison is made by plotting the addition of the catch of 1 to 6 year olds (Table 3) against the catch of bedlamers from 1952 to 1975 as calculated by Øritsland and Benjaminsen (MS 1975). The functional regression (Ricker 1973) through these points has a slope not significantly different from unity and a position not significantly different from the origin. As previously noted, the best estimates from cohort analysis are between 1961 and 1975 since during this period the catch data are more consistent. Because the estimates of population improve as the analysis proceeds (Pope 1972), 1961 should then give the most reliable abundance level. With this in mind, a projection starting in 1961, subtracting the age specific catch of the various fisheries should give the most reliable pup production, following the application of a maturity ogive. These pup productions, resulting from determining the breeding population, are probably more reliable than those predicted from sequential analysis. However, the number of animals in age groups older than one should be equivalent to the sequential analysis estimates. Recruitments to the 2+ populations were the cohort analysis abundance estimates for one year olds.

The sex ratio of the catch was distributed over ages in accordance with the asymptotic function appearing in Figure 1. The

value of the asymptote is represented by the fraction of males in the catch (Table 6). For two-year-olds, the percent of females remained 52.8%, however, the asymptotic ratio varied in relation to the closing dates of the seal hunt.

In 1961 the sex ratio was assumed to be 50:50 in the population; changing thereafter in response to the sex ratio of the hunt on the moulting lairs and the contribution of this hunt to the total catch of 1+ seals (Table 5). This assumption is reasonable since the number of adult females taken by large vessels in the breeding lairs during the postwar period was small because the value of whitecoat pelts was 2 to 6 times that of an adult pelt. Annual prices for the Norwegian fishery are given by Fiskeridirektøren (1951-1965). Prior to this time Coleman (1938) suggests that few adult females were taken in the steamboat fishery since the cost of powder and shot was in excess of the value of the pelt. During the period from 1895 to 1923 adult seals only comprised 2.7% of the total steamboat catch (Chafe et al. 1923). Even less of this would be females. From 1953 to 1960 white-coats were worth from 1.3 to 2 times as much as the adults, however, this declined to 0.7 by 1964. Furthermore, due to the labour associated with skinning and transporting the adult pelts it is unreasonable to assume that there was excessive killing of breeding females when white-coats were available.

To protect the mature females a closing date of May 5 was established in the Gulf and on the Front in 1961. At this time we assume the sex ratio in the large vessel catch was 55% adult males since the sex ratio in the population would probably be altered in favour of

females. Since the proportion of moulting females on the ice would be less than males, the fraction of females in the catch would also have been somewhat smaller than the fraction of males. In 1963 the closing date was changed to April 30. The closing date in 1965 in the Gulf was altered to April 25, and this date was also established on the front in 1968. In 1970 the closing date was April 29 and since 1971 it has been April 24. An agreement was made in 1965 that no females could be killed while breeding.

These regulations obviously decreased the percentage of mature females in the catches. The total Norwegian sample of moulting harps taken from 1969 to 1974 (Benjaminsen and Øritsland, MS 1975) showed a marked surplus of males. In age groups one and two the sex ratio is approximately 50:50 and thereafter increases to 86% males for mature animals (Fig. 1).

The projection is broken out into males and females and each component is handled separately in the simulation (Table 6). The fertility rate was assumed to be about 94% during this period, and 6% of the breeding population was assumed to be over the age of 25 (Benjaminsen and Øritsland MS 1975). A constant, conservative maturity ogive, calculated from Sergeant's (1966, 1976) data was applied to the breeding population using the following schedule:

| Age | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----------|---|-------|-------|-------|-------|-------|-------|-------|-------|
| Fraction | | | | | | | | | |
| whelping | 0 | 0.075 | 0.220 | 0.450 | 0.700 | 0.880 | 0.970 | 0.990 | 1.000 |

The sex ratio in the breeding population became as high as 54.9% in favour of females in 1971, and has continually dropped since then to 52.9% in 1976 (Table 6), as a result of a much reduced hunt for moulting animals (Table 2). Age specific sex ratios rise as seals get older due to the accumulated effect of selective hunting. In fact, in more recent years this ratio has exceeded 70% females for animals over 20 years.

The change in the sex ratio provides for more pups than would be calculated by applying a 50:50 ratio. Indeed, it is assumed that pup production calculated in this manner is more reliable than estimated from sequential analysis (Fig. 3). Production reached a minimum in 1972 of 294,071 animals, down from 424,561 produced in 1962. Since 1972 the production has slowly increased, primarily due to the entry of the 1968 year-class and the quota regulations, so that currently production would be 311,502. Using Sergeant's (1976) latest maturity estimates, production would now be approximately 330,000.

DIRECT SURVEY METHODS

Ultraviolet sensing method

Ultraviolet photography has been used for detecting certain white animals against a white background of ice or snow (Lavigne and Øritsland 1974). It is anticipated that this method could significantly improve estimates of harp seal pup production. Details of this method have been outlined by Lavigne et al. (1975), as it was applied to the northwestern Atlantic harp seal. It is stated that

using a direct statistical method i.e. applying a mean density estimate to the total herd area, results in confidence limits much too large to suggest the method is useful. In addition, the patchiness of seals on the ice results in a skewed distribution from subplot estimates such that the modal value is indeed much less than the mean.

Lavigne et al. (1975) have suggested that a ratio estimate should be used to reduce the confidence limits of the estimate. This method requires additional information over that needed for a direct estimate. Furthermore, the assumptions concerning these supplement data are critical to the estimate. In order for this method to improve upon the direct estimate the correlation of the number of adults to pups in the subplots must be greater than the ratio of the coefficient of the variance of adults among the subplots divided by 2 times the coefficient of variance of the number of pups among subplots. Lavigne et al. (1975) count the number of adults from 1220 m to get an estimate of the total herd size and determined ratio estimates at 305 m.

It is suggested that their sample size of 69 subplots was too small. However, this only required a correlation coefficient, r_{AP} , of 0.232 between pups and adults for the method to be valid. Thus the minimum assumption required for the method to be valid is

$$\frac{\sigma_A^2 \cdot P}{2\sigma_{P.A}^2} = 0.232$$

where σ^2 is the variance among subplots of pup abundance, P and

adult abundance, A. This, of course, is a minimum assumption and is subject to change depending on how good the actual agreement between the number of pups and adults. This method although perhaps the most promising of all, requires good ground truthing and a more thorough knowledge of the fraction of males present in the breeding lairs. Until these objectives are achieved it cannot be considered an adequate method, useful in the evaluation of status of the harp seal stock.

Curran's survey method

Currently, there is only one method of direct estimation of pup production which a majority of scientists agree upon as being reliable (Fig. 3). Perhaps it is more because of the man, and his long years of experience with seal management, than with the technique.

The following is a brief description of the method. On March 8th and 9th a grid is flown from northern Labrador to Notre Dame Bay, Newfoundland to locate the herd. At this time, the younger females which have less control of parturation than older seals, haul up on the ice to give birth. The sighting of these younger animals gives the herd's location. Approximately two days later the older females will begin to whelp in a somewhat more southerly location. This usually gives the appearance of two herds. The majority of the front herd have whelped by March 12th (Curran personal communication).

Each of these two herds are surveyed separately. First, the extensiveness of a herd is determined by circling it and drawing a parallel grid line. This grid is followed by aircraft and X's are

marked on it where there are no seals so the total area of seal density can be calculated.

The pups in specific subpatches are then sometimes counted by Curran and his crew for at least 20 acres. Each subpatch is about 1 acre in size and randomly distributed. In some areas the seals are scattered and in others dense, such that a mean number/acre and an associated variance can be calculated. This density is now applied to the subherd, for which the total area has been determined to give the population size. This exercise then allows for visual estimates from the aircraft for the remaining herd. Norwegian and Canadian sealers will hunt off particular areas, and Curran receives daily counts of their take. When the hunters move out he will count the remaining seals and add this to the catch, thus arriving at total figures for different areas. By relating these figures to his own estimates he can get an idea of how accurate his original estimates were. In addition to soon learning what a specific number of pups looks like from the air, he claims his method of estimation is always conservative by at least 10%; this is possibly an overstatement.

There is no doubt the method is crude and could be improved by good aerial photography. However, because of his efforts in ground truthing his direct estimates are perhaps the most accurate available.

It has been proposed that 1/3 of the seal herd whelp in the Gulf and 2/3 whelp on the Front (Sergeant 1976). According to Curran's estimates from 1971 to date, the breeding population on the

Front is about the same size as that in the Gulf when his Front estimates are subtracted from our total estimates. Although Curran's estimates are conservative, the implications of this fact are quite serious. It means that with the Gulf closed to hunting from large vessels, the Front could become severely overexploited. Apparently, the juvenile seals intermix between the two areas (Sergeant 1976) but adults do not, and therefore there is a possibility the assumption of the two herds may be one (Sergeant 1965) is wrong.

Construction of simulation

It is our opinion that some further insights can be gained into the population dynamics of harp seals by the construction of a stochastic simulation. By stochastic it is meant that the distribution and variance of certain parameters and state variables is taken into account. Another assumption is that the Front and Gulf herds are indeed one population since they spend the summer together in the Arctic (Sergeant 1965), and intermix as juveniles (Sergeant 1975). Of the two assumptions possible, more evidence supports the one herd hypothesis.

The simulations written in APL (Fig. 6) are represented by the schematic flow chart (Fig. 7). The program presented in Figure 6 requires specific large vessel catches of 1+ seals and pups to be entered however, the alternative program is slightly changed so fishing mortality rates rather than large vessel catches are entered.

To initiate the program a starting population of animals aged 1 to 25 is entered, then the number of years the program is to run,

and the starting date. A quota can then be set by entering in a large vessel catch of 1+ seals and pups. An option is also available whereby the catch of 1+ seals and pups by the Magdellanot can be fixed.

The simulation has two options related to its printout. If the number of runs is deemed as one, it will print out the breeding and total population size, the annual sex ratio of the total and breeding population, the total catch of 1+ seals and pups. However, since the model is stochastic, each run will give different answers since the "seed" in the random number generator is not fixed for each iteration. Thus, when the number of runs is greater than one, matrices of total annual and breeding population, in addition to the catch of 1+ seals and pups is printed out. The right hand two columns in each of the matrices is the mean and standard deviation for that year.

The starting population is now broken up into the male and female fractions present in 1977 (Fig. 6, [34]*). One of the problems with this fishery which does not allow it to be interpreted accurately using conventional fisheries models is that the frequency distribution of the catch bears little relationship to the frequency distribution of the population. Therefore, the catch cannot be distributed in a similar structure to the population. The frequency distributions of four district fisheries (Fig. 8) are used; the high Arctic, Greenland, landsmen and large vessel catches. These distributions are averages over a number of randomly chosen years between 1952 and 1975 so that biases due to trends in recruitment and effort are reduced. The interpolated values from the

* [n] represents the line in the program in (Fig. 6).

curves of best fit were entered into the simulation as average constant values representing the catch frequencies (Fig. 6 [45] - [52]).

All the uncontrolled catches, i.e. those excluding large vessels, had a normally distributed random component. The means and standard deviations for these fisheries were as follows:

| | Catch of 1+ seals | | |
|----------------|-------------------|------------|-----------------|
| Landsmen catch | 13026 | \pm 5048 | ISD |
| Greenland | 3784 | \pm 1040 | (Kapel 1975) |
| High Arctic | 1294 | \pm 729 | (Sergeant 1971) |

| | Catch of 0-group seals | | |
|-----------|------------------------|-------------|-----|
| Landsmen | 36949 | \pm 14442 | ISD |
| Greenland | 3784 | \pm 1040 | |

In each case, the catch was broken into the frequency distribution through multiplying it by the average catch distribution. It was then fractioned into males and females by (1) assuming that each uncontrolled catch had the same sex ratio as the current population and (2) that the large vessel catch had the same age dependent sex ratio as that in Figure 1 since the closing dates are to remain fixed.

The next step in the program is to calculate the size of the breeding population ([84]) and for this two functions must be evaluated. First density-dependent age specific whelping ages are calculated using the equation ([73]):

$$(I) \quad \text{ARC sine } {}_i E_t = 15.522A - 2.245 \times 10^{-5} N_t - 16.017$$

where ${}_i E_t$ is the fraction of the population whelping assuming 100% pregnancy for a particular age i and population number of 1+ seals N_t in the year t , and A is the age in years. The ARC sine of ${}_i E_t$ is constrained such that it cannot be 90 or 0.

A density-dependent fertility rate is calculated using the following equation ([73]):

$$(II) \quad P = 1.048 - 9.746 \times 10^{-8} N_2$$

where P is the pregnancy rate and N_2 is the number of harp seals between the ages 2 and 25. These two equations act together to constrain the pup production with the carrying capacity of the environment in addition to augmenting the production at lower population levels. They make the birth rate a power function of population size. The summation of the breeding population vector is multiplied by 1.06 since it was determined that on average 6% of the breeding population is over the age of 25.

Equation I and II represent the feedback within the simulation which changes its nature from linear to non-linear, this providing some additional realism.

Following the calculation of pup production the catches of 1+ males and females are subtracted from the total number ([86], [87]). Similarly, the pup catch is subtracted from the total catch ([94]).

Natural mortality is also considered a stochastic, normally distributed parameter of 0.114 ± 0.0302 which means that it can vary as widely as from 0.174 to 0.0536, less than 5% of the time. Natural mortality is applied using the following equation ([97] - [99]):

$$(III) \quad N_{t+1} \leftarrow N_t \text{ EXP}(-M)$$

Natural mortality is applied to the population after the catch has been subtracted off, since the hunt primarily occurs during the spring and natural mortality and hunting mortality occur quite separately.

The surviving pups after exploitation and natural mortality are assumed to have a 50:50 sex ratio. The numbers of males and females at age are now updated and the 0-group cohort is catenated into the vector for 1-year-olds and the remaining 25-year-old seals are dropped from the vector ([101], [102]). Annual sex ratios of the total and breeding populations are calculated. A test is made to determine if the simulated time period has expired and if more iterations are to occur. If the simulation time has elapsed the remaining portion of the program dealing with the calculation of means and variances and formatting executes.

Results and discussion of the simulation

Allen (1975) has pointed out that his linear model allows the population to increase ad infinitum which restricts its usefulness

for making long term predictions. Thus the initial use of the simulation was to investigate what the implications of different assumption concerning the pregnancy rates and maturity rates had on the population, when there was no fishery other than the uncontrolled landmen and aboriginal hunt.

Many mammalian populations show a varying fertility rate (Gambell 1973, Markgren 1969) in response to space and food availability, and indeed there is evidence to indicate that the pregnancy rate for harp seals can also be altered in response to changes in density (Fig. 5). When this relationship was incorporated into the simulation, it limited the population (Fig. 9) to about 6.5 million animals. At this time the fertility rate was approximately 0.4 which brought the population into equilibrium with the uncontrolled landmen and aboriginal hunt. However, the reproductive potential of the stock was unrealistic, since when MSY (maximum sustainable yield), to be discussed later, was determined, catch levels could be sustained which were in excess of those which have led to a decline in the herd.

It was apparent from this exercise that an additional mechanism was necessary to constrain the reproductive potential of the population. For this reason, a density-dependent maturity ogive was added into the simulation. The effect was to produce a population size which was in equilibrium with the landmen and aboriginal catch levels at an abundance of 3.7 million seals. The sequential analysis estimate for 1952 (Table 5) was ~2.3 million seals, which Sergeant (1975) concluded was near the maximum population size. Evidence from this

analysis do not agree with Sergeant's conclusions.

A simulated "recruitment" curve (Fig. 10) was compared with values from the sequential population analysis. The line passed through the axis of the data, but the scatter in the pup production values was too great to suggest if the two techniques confirmed one another. However, one apparent fact was that pup production had to be in excess of the 1952 production level, to sustain the high catches between 1980 and 1923 (Chafe et al. 1923).

The curve (Fig. 10) is a power function of population size. Allen (1975) speculates as to the possible shape of the recruitment curve and concludes that if reproduction is a linear function of stock size, the Beverton and Holt (1957) recruitment curve is indeed the best representation of the recruitment process of seals. In addition, to the linear shift in maturity our model allows for a linear shift in the pregnancy rate which adds more curvature to the relationship. Ricker (1954) recruitment is not realistic for a stock as undynamic as harp seals, since the declining portion of this curve, results from a population being much further out of equilibrium with the carrying capacity of the environment, than it is possible for seals to get. At the point of maximum population size then, the reproductive rate is equivalent to the mortality rate.

It was necessary to determine the MSY population level as a reference point for harp seal management. This could not be determined from the simulation if only catch was controlled, since under these conditions the population could only come into equilibrium or collapse.

Therefore, the alternate program was utilized such that catch would vary in response to population size, by controlling the hunting mortality rate. This allows for a Shaefer type curve (Fig. 11) to be generated. Hunting mortality by large vessels was calculated under present conditions to be approximately 0.01 on 1+ seals. Holding this mortality constant, the hunting level on pups was allowed to vary. The predicted value for maximum sustainable yield approximately 200,000 pups and 40,000 1+ seals, with respect to the present pup to 1+ seals kill ratio. The MSY population size is near 1.6 million 1+ seals, or a breeding stock of 375,000 females.

The variance in catch becomes greater on the left hand side of the Shaefer type curve (Fig. 11). This fact has been speculated on by Doubleday (1976), but here the biological basis is apparent. At stock sizes less than 1.2 million, the maturity ogive can no longer shift to the left thus the population loses much of its density dependent control to maintain stability. This same result was shown by Lett and Kohler (1976) for an Atlantic herring stock.

Using the 1977 age specific abundance levels (Table 5) the population was projected ahead with fixed large vessel catches of pups and 1+ seals (Fig. 12). When the large vessels removed 10,000 1+ seals and 80,000 pups the mean population size increased, reaching the MSY breeding stock size by 1968, however, there is a 66% chance MSY could not be reached until 1983. The total average catch including landmen, large vessels, Canadian native and Greenland is ~150,000 seals. When the large vessels take 10,000 1+ seals and 100,000 pups the mean population

size reaches MSY in 1991, a date which is not much different than that for the projected 1971-1974 management strategy. In all iterations the population increased at this level of exploitation.

When the large vessel catch was 10,000 1+ seals and 120,000 pups for a total average catch of 190,000, the population did not change in size. However, 20% of the time the population declined and 80% of the time it stayed the same or increased. When 10,000 more 1+ seals were removed by large vessels (Fig. 12D) the population declined in all cases.

In conclusion, this study would indicate that the catch should not exceed 170,000 seals to allow the population to increase to MSY, assuming kill ratio of 20% 1+ seals to 80% pups and keeping in mind this total catch includes the high Arctic and Greenland hunt. Furthermore, because of the growing uncertainties of prediction encountered as one moves away from the current population size (Fig. 12) quotas should not be set more than 3-5 years in advance. A complete re-examination of the population dynamics and herd assessment is necessary at least every 5 years.

Sampling of the landmen and large vessel catch must be continued at an accelerated level in the interim, in addition to samples collected from the moulting lairs for estimates of natural mortality. Furthermore, samples of ovaries should be collected from the breeding females to detect shifts in the maturity ogive. Ultraviolet aerial sensing of the herd, with adequate ground truthing, must be pursued until a direct estimate of the population size is available. We believe that the

model presented in this paper is quite complete as far as the population dynamics is concerned, and for this reason a logical extension of the simulation could incorporate some community structure, and bioenergetic submodels.

Acknowledgement

We would like to thank all those who collected the samples over the years and crewed the research vessels. In particular, we express our appreciation to D. M. Lavigne, T. Øritsland and D. E. Sergeant for allowing us to use their unpublished data. B. Beck and W. Hoek assisted in the data reduction from the files of the Arctic Biological Station.

Throughout the construction of the model and writing of this paper, T. Curran benevolently passed on much of his knowledge concerning the management of the herd during many informal conversations. P.F.L. greatly appreciates the help in data analysis through consultation with D. M. Lavigne during some initial attempts to construct a realistic population model for this herd. No doubt some of those original ideas have found their way into this present work.

Many individuals have reviewed data and ideas over the past two years pertinent to the 'genesis' of this paper, in addition to thoroughly reviewing the model as it now stands. Therefore we thank, B. Bergflødt, C. Capstick, D. Fisher, Sv. A. Horsted, F. Kapel, D. M. Lavigne, A. Mansfield, T. Øritsland, W. E. Ricker, K. R. Ronald, D. E. Sergeant, and Ø. Ulltang. C. Capstick in particular, was very

helpful in his honest appraisal of the assumptions in this paper, and the reprogramming of these assumptions in his model, to check on the outputs from the simulation now presented.

P.F.L. accepts any responsibility connected with incorrect conclusions resulting from errors in the many computer programs necessary to analyse the data and build the simulations.

- Allen, R. L. 1975. A life table for harp seals (*Pagophilus groenlandicus* (Erxleben). Symposium on the Biology of the Seal, Guelph, Canada, August 1972. Rapp. P-v Réun. Cons. per. int. Explor. Mer. 1969: 303-311.
- Benjaminsen, T. and T. Øritsland. MS 1975. Adjusted estimates of year-class survival and production with estimates of mortality for Northwestern Atlantic harp seals. Int. Comm. Northwest Atl. Fish. Working Pap., 1975(XI/3): 38 p.
- Beverton, R. J. H., and S. J. Holt. 1957. On the dynamics of exploited fish populations. Fishery Investigations, Ser. II, Vol. 19, 533 p.
- Bourlière, F. 1959. Lifespans of mammalian and bird populations in nature. In G.E.W. Wolstenholme and Maeve O'Connor [ed] Colloquia on Ageing. J. and A. Churchill Ltd., London.
- Capstick, C. K., and K. Ronald. 1976. Modelling seal populations for herd management. FAO Scientific Consultation on Marine Mammals, Bergen, Norway. ACMRR/MM/SC/77. 19 p.
- Chafe, L. G., W. A. Mann, and H. M. Mosdell. 1923. Chafe's Sealing Book, a history of the Newfoundland sealfishery from the earliest available records down to and including the voyage of 1923. The Trade Printers and Publishers Ltd. 108 p.
- Coleman, J. 1938. The Newfoundland seal fishery and its possible influence on the Greenlanders. Polar Record, 2(16): 99-103.
- Doubleday, W. G. 1975. The least squares approach to analysing catch at age data. Int. Comm. Northwest Atl. Fish. Res. Doc. 75/35.

- Doubleday, W. G. 1976. Environmental fluctuations and fisheries management. Int. Comm. Northwest. Atl. Fish. Sel. Pap. 1.1: 141-150.
- Fisher, H. D. 1952. Harp seals of the Northwest Atlantic. Fish. Res. Board Can. General Ser. No. 20, 4 p.
- Fisher, H. D. 1955. Utilization of Atlantic harp seal populations. Trans. 20 North American Wildlife Conference: 507-518.
- Fiskeridirektøren, 1951 - 1965. Selfangsten Arsberetn. Norg. Fisk. 1946-1964.
- Fry, F. E. J. 1949. Statistics of a lake trout fishery. Biometrics 5: 27-67.
- Gambell, R. 1973. Some effects of exploitation on reproduction in whales. J. Reprod. Fert., Suppl. 19: 533-553.
- Gulland, J. A. 1965. Estimation of mortality rates. Annex to Arctic Fisheries Working Group Report. ICES C.M. 1965 (3).
- International Commission for the Northwest Atlantic Fisheries 1972. Int. Comm. Northwest Atl. Fish. Sim. Doc. 72/6.
- Jones, R., 1964. Estimating population size from commercial statistics when fishing mortality varies with age. Rapp. P.v. Cons. perm. int. explor. Mer. 155,38: 210-214.
- Kapel, F. O. 1975. Data on the catch of harp and hooded seals 1954-1974, and long-term fluctuations in sealing in Greenland. Int. Comm. Northwest. Atl. Fish. Res. Doc. 75/XII/9.

- Lavigne, D. M., and N. A. Øritsland. 1974. Ultraviolet photography: A new method for remote sensing of mammals. *Can. J. Zool.* 52: 939-941.
- Lavigne, D. M., S. Innes, K. Kalpakis and K. Ronald. 1975. An aerial census of western Atlantic harp seals (*Pagophilus groenlandicus*) using ultraviolet photography. *Int. Comm. Northwest Atl. Fish. Res. Doc.* 75/XII/144.
- Lavigne, D. M., W. Barchard, S. Innes, and N. A. Øritsland. 1976. Pinniped bioenergetics. *FAO. Scientific Consultation on Marine Mammals. Bergen, Norway.* ACMRR/MM/SC/112.
- Lett, P. F. and A. C. Kohler. 1976. Recruitment: a problem of multi-species interaction and environmental perturbations, with special reference to Gulf of St. Lawrence Atlantic herring (*Clupea harengus harengus*). *J. Fish. Res. Board Can* 33: 1353-1371.
- Markgren, G. 1969. Reproduction of moose in Sweden. *Viltrevy* 6 (3): 125-299.
- McLaren, I. A. 1967. Seals and group selection. *Ecology* 48(1): 104-110.
- Murphy, G.I. 1964. A solution of the catch equation. *J. Fish. Res. Board Can.* 22: 191-201.
- Øritsland, T. 1967. Revised catch and effort statistics for the Norwegian seal hunt in the Front and Gulf areas off Newfoundland, 1937-1967. *Int. Comm. Northwest Atl. Fish. Serial No.* 1959.

- Øritsland, T. 1970. Sealing and seal research in the southwest Atlantic pack ice, Sept.-Oct. 1964, p. 367-376. In Antarctic ecology [ed] M. W. Hollgate Vol. 1. Academic Press, London.
- Øritsland, T. 1971. Catch and effort statistics for Norwegian sealing in Front area, Newfoundland, 1971. Int. Comm. Northwest Atl. Fish. Res. Doc. 71/132.
- Øritsland, T., and T. Benjaminsen. MS 1975. Further documentation of comments of Canadian harp seal estimates. Int. Comm. Northwest Atl. Fish. Working Pap., 1975 (XII/7): 1-9.
- Pope, J. G. 1972. An investigation of the accuracy of virtual population analysis using cohort analysis. Int. Comm. Northwest Atl. Fish. Res. Bull. 9: 65-74.
- Ricker, W. E. 1954. Stock and recruitment. J. Fish. Res. Board Can. 11: 559-623.
- Ricker, W. E. 1971. Comments on the West Atlantic harp seal herd and proposals for the 1972 harvest. Int. Comm. Northwest Atl. Fish. Res. Doc. 75/IXX/143 (Appendix).
- Ricker, W. E. 1973. Linear regressions in fishery research. J. Fish. Res. Bd Can., 30: 409-434.
- Sergeant, D. E. 1965. Migration of harp seal *Pagophilus groenlandicus* in the northwest Atlantic. J. Fish. Res. Board Can. 22: 433-464.
- Sergeant, D. E. 1966. Reproductive rates of harp seals, *Pagophilus groenlandicus* (Erxleben). J. Fish. Res. Board Can. 23: 757-766.

- Sergeant, D. E. 1969. On the population dynamics and size of stocks of harp seals in the northwest Atlantic. Int. Comm. Northwest Atl. Fish. Res. Doc. 69/31: 1-25.
- Sergeant, D. E. 1970. Results of research on harp seals in 1969-70. Int. Comm. Northwest Atl. Fish. Res. Doc. 70/61: 1-15.
- Sergeant, D. E. 1971. Calculation of production of harp seals in the western north Atlantic. Int. Comm. Northwest Atl. Fish. Redbook 1971, Part III: 157-184.
- Sergeant, D. E. 1972. Canadian studies on harp seals in 1971. Int. Comm. Northwest Atl. Fish. Res. Doc. 72/59.
- Sergeant, D. E. 1973a. Feeding, growth and productivity of northwest Atlantic harp seals (*Pagophilus groenlandicus*). J. Fish. Res. Bd. Can. 30: 17-29.
- Sergeant, D. E. 1973b. Environment and reproduction in seals. J. Reprod. Fert., Suppl. 19: 555-561.
- Sergeant, D. E. 1975. Results of research on harp seals in 1975 with an estimate of production. Int. Comm. Northwest. Atl. Fish. Res. Doc. 75/XII/142.
- Sergeant, D. E. 1976. Studies of harp seals of the western north Atlantic populations in 1976. Int. Comm. Northwest. Atl. Fish. Res. Doc. 76/X/124.
- Sergeant, D. E. and H. D. Fisher. 1960. Harp seal populations in the western North Atlantic from 1950 to 1960. Fish. Res. Board Can. Circular No. 5, 58 p.

Simms, H. S., B. N. Berg, and D. F. Davies. 1959. Onset of disease and the longevity of rat and man. In G. E. W. Wolstenholme and Maeve O'Connor [ed] Colloquia on Ageing. J. and A. Churchill Ltd. London.

Ulltang, Ø. MS 1971. Estimates of mortality and production of harp seals at Newfoundland. Working Pap. Int. Comm. Northwest Atl. Fish. Special Meet. Panel A Experts, Charlottenlund, Denmark, 23-24 Sept. 1971. 13 p.

Table 1. Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel A | Landsmen A | % 1952 | Large vessel A | Landsmen A | % 1953 | Large vessel A | Landsmen A | % 1954 | Large vessel Avg. | Landsmen A | % 1955 |
|-----|-------------------|---------------|-----------|-------------------|---------------|-----------|-------------------|---------------|-----------|----------------------|---------------|-----------|
| 1 | 10 | 27 | 4.29 | 419 | 15 | 27.61 | 233 | 15 | 35.76 | 30.71% | 48 | 29.28 |
| 2 | 24 | 95 | 10.57 | 124 | 52 | 8.71 | 91 | 26 | 14.28 | 13.99 | 77 | 14.07 |
| 3 | 15 | 98 | 6.96 | 103 | 98 | 7.86 | 26 | 23 | 4.31 | 7.70 | 98 | 8.48 |
| 4 | 11 | 114 | 5.49 | 61 | 108 | 5.22 | 36 | 57 | 6.52 | 6.02 | 41 | 6.16 |
| 5 | 13 | 115 | 6.30 | 62 | 111 | 5.32 | 19 | 45 | 3.63 | 4.90 | 58 | 5.34 |
| 6 | 25 | 101 | 11.03 | 48 | 100 | 4.28 | 24 | 33 | 4.26 | 4.20 | 35 | 4.38 |
| 7 | 17 | 96 | 7.75 | 43 | 86 | 3.80 | 21 | 40 | 3.92 | 3.64 | 30 | 3.79 |
| 8 | 17 | 81 | 7.61 | 48 | 51 | 3.72 | 20 | 23 | 3.39 | 3.28 | 39 | 3.57 |
| 9 | 12 | 60 | 5.40 | 47 | 50 | 3.65 | 13 | 16 | 2.27 | 2.94 | 13 | 2.91 |
| 10 | 14 | 54 | 6.15 | 39 | 41 | 3.02 | 21 | 10 | 3.31 | 2.72 | 14 | 2.72 |
| 11 | 12 | 60 | 5.40 | 51 | 28 | 3.66 | 8 | 8 | 1.37 | 2.46 | 10 | 2.43 |
| 12 | 3 | 42 | 1.60 | 28 | 34 | 2.22 | 18 | 7 | 2.87 | 2.27 | 13 | 2.29 |
| 13 | 3 | 21 | 1.41 | 22 | 22 | 1.69 | 18 | 5 | 2.84 | 2.10 | 2 | 1.99 |
| 14 | 5 | 12 | 2.13 | 17 | 19 | 1.33 | 9 | 4 | 1.45 | 1.96 | 3 | 1.87 |
| 15 | 9 | 24 | 3.86 | 33 | 34 | 2.55 | 12 | 4 | 1.90 | 1.82 | 5 | 1.77 |
| 16 | 3 | 12 | 1.32 | 37 | 10 | 2.54 | 16 | 0 | 2.44 | 1.74 | 4 | 1.67 |
| 17 | 5 | 15 | 2.16 | 26 | 14 | 1.86 | 7 | 2 | 1.11 | 1.54 | 3 | 1.48 |
| 18 | 4 | 8 | 1.69 | 16 | 11 | 1.17 | 2 | 5 | 0.40 | 1.34 | 4 | 1.31 |
| 19 | 2 | 8 | 0.88 | 12 | 6 | 0.85 | 8 | 2 | 1.26 | 1.26 | 2 | 1.21 |
| 20 | 11 | 12 | 4.56 | 50 | 39 | 3.72 | 4 | 8 | 0.76 | 1.01 | 2 | 0.97 |
| 21 | 3 | 6 | 1.26 | 32 | 7 | 2.22 | 2 | 3 | 0.36 | 0.76 | 4 | 0.76 |
| 22 | 0 | 6 | 0.05 | 18 | 3 | 1.21 | 2 | 2 | 0.34 | 0.67 | 3 | 0.67 |
| 23 | 1 | 4 | 0.44 | 12 | 7 | 0.87 | 5 | 1 | 0.78 | 0.50 | | 0.47 |
| 24 | 3 | 2 | 1.23 | 9 | 3 | 0.62 | 1 | 0 | 0.15 | 0.42 | | 0.39 |
| 25 | 1 | 4 | 0.44 | 5 | 1 | 0.33 | 2 | 1 | 0.32 | 0.06 | | 0.05 |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landmen to larger vessel catch.

| Age | Large vessel | | Landmen | | % 1956 | | Large vessel | | Landmen | | % 1957 | | Large vessel | | Landmen | | % 1958 | | Large vessel | | Landmen | | % 1959 | | Large vessel | | Landmen | | % 1960 | |
|-----|--------------|---|---------|---|--------|---|--------------|---|---------|---|--------|---|--------------|---|---------|---|--------|---|--------------|---|---------|---|--------|---|--------------|---|---------|---|--------|--|
| | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | A | |
| 1 | | | 11 | | 27.42 | | 11 | | 29.78 | | 40 | | 26 | | 15.43 | | 9 | | 29.37 | | 34 | | 19.49 | | | | | | | |
| 2 | | | 59 | | 13.42 | | 20 | | 13.64 | | 15 | | 25 | | 7.14 | | 23 | | 13.74 | | 107 | | 9.12 | | | | | | | |
| 3 | | | 62 | | 7.88 | | 23 | | 7.67 | | 19 | | 90 | | 4.01 | | 25 | | 7.79 | | 67 | | 5.17 | | | | | | | |
| 4 | | | 55 | | 6.27 | | 26 | | 6.11 | | 22 | | 142 | | 3.19 | | 18 | | 6.06 | | 202 | | 4.02 | | | | | | | |
| 5 | | | 37 | | 4.97 | | 17 | | 4.91 | | 18 | | 110 | | 2.57 | | 23 | | 5.09 | | 47 | | 3.36 | | | | | | | |
| 6 | | | 35 | | 4.32 | | 15 | | 4.21 | | 11 | | 115 | | 2.21 | | 26 | | 4.48 | | 62 | | 2.97 | | | | | | | |
| 7 | | | 42 | | 3.94 | | 21 | | 3.77 | | 11 | | 82 | | 1.97 | | 25 | | 3.93 | | 41 | | 2.60 | | | | | | | |
| 8 | | | 54 | | 3.82 | | 17 | | 3.36 | | 9 | | 81 | | 1.76 | | 22 | | 3.52 | | 19 | | 2.34 | | | | | | | |
| 9 | | | 45 | | 3.37 | | 16 | | 3.03 | | 9 | | 58 | | 1.59 | | 21 | | 3.19 | | 14 | | 2.11 | | | | | | | |
| 10 | | | 33 | | 2.97 | | 21 | | 2.88 | | 17 | | 69 | | 1.51 | | 17 | | 2.90 | | 18 | | 1.92 | | | | | | | |
| 11 | | | 23 | | 2.58 | | 21 | | 2.64 | | 13 | | 63 | | 1.38 | | 10 | | 2.53 | | 13 | | 1.67 | | | | | | | |
| 12 | | | 30 | | 2.52 | | 18 | | 2.42 | | 14 | | 45 | | 1.26 | | 7 | | 2.29 | | 12 | | 1.52 | | | | | | | |
| 13 | | | 19 | | 2.19 | | 12 | | 2.18 | | 8 | | 40 | | 1.13 | | 5 | | 2.09 | | 6 | | 1.39 | | | | | | | |
| 14 | | | 20 | | 2.07 | | 16 | | 2.09 | | 6 | | 31 | | 1.10 | | 9 | | 2.03 | | 9 | | 1.35 | | | | | | | |
| 15 | | | 11 | | 1.80 | | 14 | | 1.93 | | 16 | | 27 | | 1.01 | | 7 | | 1.86 | | 4 | | 1.23 | | | | | | | |
| 16 | | | 20 | | 1.88 | | 9 | | 1.78 | | 14 | | 14 | | 0.90 | | 4 | | 1.72 | | 5 | | 1.15 | | | | | | | |
| 17 | | | 15 | | 1.62 | | 8 | | 1.58 | | 6 | | 18 | | 0.80 | | 4 | | 1.54 | | 3 | | 1.02 | | | | | | | |
| 18 | | | 25 | | 1.62 | | 8 | | 1.39 | | 3 | | 13 | | 0.72 | | 2 | | 1.32 | | 3 | | 0.87 | | | | | | | |
| 19 | | | 15 | | 1.37 | | 6 | | 1.29 | | 3 | | 5 | | 0.67 | | 3 | | 1.25 | | 0 | | 0.83 | | | | | | | |
| 20 | | | 16 | | 1.17 | | 11 | | 1.11 | | 9 | | 15 | | 0.58 | | 2 | | 0.99 | | 4 | | 0.66 | | | | | | | |
| 21 | | | 10 | | 0.84 | | 6 | | 0.85 | | 7 | | 3 | | 0.42 | | 0 | | 0.72 | | 0 | | 0.48 | | | | | | | |
| 22 | | | 8 | | 0.73 | | 5 | | 0.71 | | 6 | | 3 | | 0.37 | | 0 | | 0.64 | | 1 | | 0.42 | | | | | | | |
| 23 | | | 9 | | 0.60 | | 0 | | 0.48 | | 2 | | 2 | | 0.25 | | 1 | | 0.50 | | 0 | | 0.33 | | | | | | | |
| 24 | | | 7 | | 0.49 | | 2 | | 0.43 | | 4 | | 4 | | 0.22 | | 0 | | 0.40 | | 0 | | 0.27 | | | | | | | |
| 25 | | | 3 | | 0.10 | | 3 | | 0.09 | | 2 | | 2 | | 0.05 | | 1 | | 0.07 | | 0 | | 0.05 | | | | | | | |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | | Landsman | | % 1961 | | Large vessel | | Landsman | | % 1962 | | Large vessel | | Landsman | | % 1963 | | Large vessel | | Landsman | | % 1964 | |
|-----|--------------|-----|----------|----|--------|-----|--------------|-----|----------|----|--------|-------|--------------|----|----------|-----|--------|---|--------------|---|----------|---|--------|---|
| | A | | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B |
| 1 | 122 | 16 | 26.35 | 20 | 6 | 26 | 23.55 | 111 | 18 | 26 | 44 | 7.93 | 3 | 10 | 43 | 53 | 3.13 | | | | | | | |
| 2 | 34 | 65 | 10.39 | 50 | 15 | 65 | 29.17 | 121 | 52 | 83 | 135 | 10.28 | 6 | 31 | 42 | 73 | 5.53 | | | | | | | |
| 3 | 25 | 100 | 10.27 | 74 | 16 | 90 | 8.25 | 106 | 86 | 54 | 140 | 9.42 | 8 | 74 | 50 | 124 | 7.97 | | | | | | | |
| 4 | 30 | 126 | 12.62 | 95 | 30 | 125 | 8.15 | 55 | 90 | 33 | 123 | 5.83 | 11 | 64 | 32 | 96 | 9.30 | | | | | | | |
| 5 | 11 | 39 | 4.27 | 39 | 21 | 116 | 6.07 | 46 | 80 | 26 | 106 | 4.93 | 7 | 65 | 32 | 97 | 6.72 | | | | | | | |
| 6 | 13 | 52 | 5.34 | 33 | 7 | 40 | 2.60 | 61 | 56 | 21 | 77 | 5.35 | 12 | 40 | 42 | 82 | 9.65 | | | | | | | |
| 7 | 15 | 43 | 5.30 | 46 | 14 | 60 | 2.54 | 67 | 32 | 8 | 40 | 5.04 | 6 | 27 | 29 | 56 | 5.15 | | | | | | | |
| 8 | 9 | 27 | 3.24 | 16 | 16 | 58 | 2.63 | 66 | 36 | 10 | 46 | 5.09 | 5 | 11 | 35 | 46 | 4.28 | | | | | | | |
| 9 | 8 | 28 | 3.08 | 17 | 13 | 42 | 2.53 | 63 | 22 | 12 | 34 | 4.67 | 5 | 10 | 23 | 33 | 4.00 | | | | | | | |
| 10 | 17 | 24 | 4.76 | 18 | 16 | 34 | 1.15 | 72 | 25 | 15 | 40 | 5.36 | 8 | 6 | 27 | 33 | 5.95 | | | | | | | |
| 11 | 10 | 15 | 2.84 | 14 | 10 | 24 | 1.24 | 70 | 24 | 14 | 38 | 5.20 | 4 | 11 | 18 | 29 | 3.25 | | | | | | | |
| 12 | 3 | 20 | 1.63 | 17 | 3 | 20 | 1.77 | 51 | 20 | 10 | 30 | 3.83 | 4 | 5 | 21 | 26 | 3.19 | | | | | | | |
| 13 | 4 | 16 | 1.64 | 11 | 11 | 22 | 0.99 | 57 | 11 | 9 | 20 | 4.03 | 3 | 6 | 20 | 26 | 2.53 | | | | | | | |
| 14 | 6 | 7 | 1.61 | 9 | 13 | 21 | 1.32 | 56 | 17 | 14 | 31 | 4.17 | 3 | 3 | 21 | 24 | 2.48 | | | | | | | |
| 15 | 1 | 12 | 0.81 | 12 | 23 | 44 | 1.98 | 42 | 22 | 11 | 33 | 3.31 | 5 | 4 | 7 | 11 | 3.50 | | | | | | | |
| 16 | 5 | 5 | 1.29 | 7 | 10 | 17 | 0.69 | 54 | 15 | 9 | 24 | 3.91 | 4 | 2 | 11 | 13 | 2.90 | | | | | | | |
| 17 | 4 | 3 | 0.98 | 12 | 9 | 13 | 1.56 | 36 | 13 | 11 | 24 | 2.76 | 5 | 4 | 12 | 16 | 3.62 | | | | | | | |
| 18 | 2 | 2 | 0.52 | 5 | 4 | 15 | 0.78 | 24 | 9 | 4 | 13 | 1.78 | 8 | 1 | 6 | 7 | 5.37 | | | | | | | |
| 19 | 3 | 0 | 0.63 | 4 | 6 | 7 | 0.56 | 18 | 4 | 12 | 16 | 1.45 | 4 | 0 | 14 | 14 | 2.92 | | | | | | | |
| 20 | 2 | 7 | 0.77 | 9 | 4 | 16 | 1.26 | 18 | 10 | 10 | 20 | 1.53 | 0 | 2 | 6 | 8 | 0.18 | | | | | | | |
| 21 | 0 | 0 | 0.21 | 1 | 2 | 3 | 0.16 | 17 | 5 | 6 | 11 | 1.30 | 4 | 0 | 6 | 6 | 2.74 | | | | | | | |
| 22 | 2 | 0 | 0.42 | 4 | 5 | 11 | 0.61 | 12 | 5 | 8 | 13 | 1.01 | 2 | 2 | 3 | 5 | 1.42 | | | | | | | |
| 23 | 1 | 2 | 0.31 | 0 | 1 | 4 | 0.17 | 9 | 4 | 10 | 14 | 0.84 | 2 | 0 | 5 | 5 | 1.42 | | | | | | | |
| 24 | 0 | 1 | 0.26 | 1 | 2 | 3 | 0.16 | 8 | 2 | 6 | 8 | 0.66 | 3 | 1 | 2 | 3 | 2.02 | | | | | | | |
| 25 | 1 | 4 | 0.41 | 1 | 2 | 5 | 0.18 | 4 | 1 | 2 | 3 | 0.31 | 1 | 0 | 6 | 6 | 0.79 | | | | | | | |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | | Landsman | | Total | | 1965 | | 1966 | | Large vessel | | Landsman | | Total | | 1967 | |
|-----|--------------|----|----------|-----|-------|----|------|----|-------|-----|--------------|-----|----------|----|-------|-----|-------|---|
| | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B | A | B |
| 1 | 32 | 4 | 4 | 4 | 120 | 10 | 1 | 11 | 15.30 | 176 | 55 | 231 | 17 | 12 | 6 | 35 | 18.53 | |
| 2 | 13 | 17 | 10 | 10 | 100 | 29 | 2 | 31 | 13.75 | 61 | 31 | 92 | 14 | 42 | 11 | 67 | 8.68 | |
| 3 | 13 | 45 | 24 | 69 | 39 | 42 | 3 | 45 | 6.86 | 18 | 5 | 23 | 1 | 46 | 17 | 64 | 3.34 | |
| 4 | 15 | 72 | 34 | 106 | 40 | 44 | 5 | 49 | 7.17 | 18 | 6 | 24 | 9 | 37 | 24 | 70 | 3.56 | |
| 5 | 23 | 71 | 24 | 95 | 45 | 42 | 7 | 49 | 7.79 | 29 | 19 | 48 | 11 | 58 | 30 | 99 | 6.11 | |
| 6 | 22 | 66 | 18 | 84 | 45 | 50 | 5 | 55 | 8.06 | 46 | 22 | 68 | 21 | 72 | 25 | 118 | 8.11 | |
| 7 | 7 | 39 | 15 | 54 | 44 | 33 | 4 | 37 | 7.15 | 40 | 19 | 59 | 26 | 64 | 27 | 117 | 7.39 | |
| 8 | 2 | 32 | 16 | 48 | 31 | 19 | 0 | 19 | 4.69 | 30 | 13 | 43 | 21 | 43 | 14 | 78 | 5.21 | |
| 9 | 2 | 18 | 4 | 22 | 15 | 14 | 4 | 18 | 2.67 | 23 | 5 | 28 | 26 | 31 | 10 | 67 | 3.79 | |
| 10 | 5 | 9 | 5 | 14 | 17 | 8 | 2 | 10 | 2.55 | 16 | 8 | 24 | 16 | 27 | 9 | 52 | 3.12 | |
| 11 | 1 | 10 | 5 | 15 | 23 | 10 | 1 | 11 | 3.34 | 34 | 13 | 36 | 21 | 14 | 11 | 46 | 3.89 | |
| 12 | 7 | 12 | 6 | 18 | 17 | 5 | 0 | 5 | 2.32 | 18 | 6 | 24 | 14 | 15 | 5 | 34 | 2.67 | |
| 13 | 1 | 16 | 7 | 23 | 15 | 7 | 0 | 7 | 2.17 | 14 | 5 | 19 | 16 | 10 | 6 | 32 | 2.24 | |
| 14 | 5 | 8 | 7 | 15 | 17 | 7 | 0 | 7 | 2.42 | 19 | 10 | 29 | 15 | 16 | 7 | 38 | 3.15 | |
| 15 | 4 | 8 | 10 | 18 | 14 | 7 | 0 | 7 | 2.05 | 20 | 10 | 30 | 13 | 12 | 7 | 32 | 3.08 | |
| 16 | 2 | 6 | 6 | 12 | 14 | 7 | 1 | 8 | 2.09 | 14 | 6 | 20 | 17 | 4 | 6 | 27 | 2.19 | |
| 17 | 1 | 5 | 5 | 10 | 10 | 3 | 2 | 5 | 1.46 | 18 | 8 | 26 | 12 | 9 | 3 | 24 | 3.34 | |
| 18 | 3 | 2 | 4 | 6 | 17 | 1 | 0 | 1 | 2.14 | 20 | 9 | 29 | 9 | 7 | 4 | 20 | 2.71 | |
| 19 | 3 | 6 | 4 | 10 | 10 | 1 | 2 | 3 | 1.37 | 21 | 9 | 30 | 8 | 4 | 6 | 18 | 2.74 | |
| 20 | 1 | 3 | 1 | 4 | 11 | 2 | 0 | 2 | 1.45 | 15 | 4 | 19 | 11 | 5 | 4 | 20 | 1.95 | |
| 21 | 2 | 3 | 2 | 5 | 8 | 1 | 0 | 1 | 1.03 | 8 | 4 | 12 | 3 | 3 | 4 | 10 | 1.16 | |
| 22 | 1 | 1 | 1 | 2 | 3 | 1 | 1 | 2 | 0.46 | 8 | 1 | 9 | 2 | 4 | 0 | 6 | 0.84 | |
| 23 | 0 | 0 | 2 | 2 | 7 | 1 | 0 | 1 | 0.91 | 8 | 3 | 11 | 3 | 0 | 1 | 4 | 0.94 | |
| 24 | 0 | 1 | 2 | 3 | 4 | 0 | 0 | 0 | 0.49 | 4 | 2 | 6 | 5 | 2 | 2 | 9 | 0.68 | |
| 25 | 0 | 1 | 2 | 3 | 2 | 0 | 2 | 2 | 0.34 | 4 | 2 | 6 | 1 | 3 | 1 | 5 | 0.58 | |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | | Landsman | | Total Landsman | % 1968 | Large vessel | | | Landsman | | | | Total Landsman | % 1969 |
|-----|--------------|-----|----------|----|-------------------|-----------|--------------|-----|-----|----------|----|----|----|-------------------|-----------|
| | A | B | A | B | | | A | B | A | B | C | D | | | |
| 1 | 84 | 142 | 7 | 11 | 18 | 15.04 | 62 | 396 | 458 | 87 | 37 | 9 | 5 | 138 | 35.77 |
| 2 | 77 | 78 | 8 | 48 | 56 | 11.37 | 2 | 56 | 58 | 13 | 14 | 20 | 4 | 51 | 5.10 |
| 3 | 48 | 43 | 8 | 68 | 76 | 7.71 | 6 | 40 | 46 | 13 | 21 | 49 | 29 | 112 | 5.26 |
| 4 | 20 | 27 | 8 | 56 | 64 | 4.58 | 1 | 32 | 33 | 13 | 10 | 69 | 18 | 110 | 4.28 |
| 5 | 17 | 32 | 5 | 39 | 44 | 4.23 | 2 | 43 | 45 | 11 | 11 | 71 | 11 | 104 | 5.05 |
| 6 | 23 | 20 | 9 | 54 | 63 | 4.30 | 4 | 30 | 34 | 10 | 2 | 49 | 6 | 67 | 3.62 |
| 7 | 27 | 48 | 12 | 41 | 53 | 6.13 | 1 | 38 | 39 | 5 | 5 | 60 | 11 | 81 | 4.22 |
| 8 | 39 | 38 | 15 | 39 | 54 | 6.28 | 6 | 45 | 51 | 7 | 4 | 59 | 10 | 80 | 5.08 |
| 9 | 23 | 31 | 16 | 17 | 33 | 4.29 | 2 | 39 | 41 | 1 | 2 | 43 | 11 | 57 | 3.96 |
| 10 | 26 | 30 | 15 | 18 | 33 | 4.42 | 0 | 29 | 29 | 6 | 11 | 40 | 2 | 59 | 3.12 |
| 11 | 19 | 23 | 10 | 14 | 24 | 3.30 | 3 | 24 | 27 | 4 | 6 | 38 | 3 | 51 | 2.84 |
| 12 | 18 | 12 | 13 | 14 | 27 | 2.59 | 1 | 18 | 19 | 4 | 2 | 22 | 6 | 34 | 1.97 |
| 13 | 16 | 16 | 9 | 7 | 16 | 2.46 | 2 | 21 | 23 | 5 | 5 | 20 | 1 | 31 | 2.21 |
| 14 | 24 | 18 | 42 | 7 | 16 | 3.10 | 2 | 24 | 26 | 3 | 2 | 16 | 1 | 22 | 2.27 |
| 15 | 20 | 15 | 35 | 9 | 19 | 2.72 | 2 | 24 | 26 | 3 | 1 | 24 | 0 | 28 | 2.37 |
| 16 | 20 | 14 | 34 | 11 | 17 | 2.61 | 2 | 16 | 18 | 2 | 1 | 18 | 1 | 22 | 1.69 |
| 17 | 11 | 17 | 28 | 10 | 19 | 2.27 | 0 | 24 | 24 | 6 | 5 | 11 | 0 | 22 | 2.13 |
| 18 | 17 | 27 | 44 | 7 | 11 | 3.11 | 0 | 21 | 21 | 2 | 0 | 11 | 1 | 14 | 1.77 |
| 19 | 16 | 22 | 38 | 5 | 10 | 2.70 | 0 | 22 | 22 | 3 | 2 | 5 | 0 | 10 | 1.78 |
| 20 | 11 | 16 | 27 | 3 | 5 | 1.87 | 0 | 20 | 20 | 2 | 0 | 4 | 1 | 7 | 1.58 |
| 21 | 6 | 11 | 17 | 4 | 1 | 1.22 | 1 | 13 | 14 | 2 | 0 | 5 | 1 | 8 | 1.16 |
| 22 | 5 | 19 | 24 | 4 | 3 | 1.72 | 2 | 10 | 12 | 1 | 1 | 2 | 0 | 4 | 0.94 |
| 23 | 4 | 10 | 14 | 2 | 0 | 0.95 | 2 | 9 | 11 | 2 | 0 | 0 | 0 | 2 | 0.84 |
| 24 | 2 | 8 | 10 | 1 | 0 | 0.67 | 1 | 5 | 6 | 0 | 0 | 1 | 0 | 1 | 0.45 |
| 25 | 1 | 4 | 5 | 0 | 1 | 0.35 | 0 | 7 | 7 | 0 | 0 | 1 | 0 | 1 | 0.53 |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | | Landsman | | | | Total Landsman | % 1970 | Large vessel | Landsman | | | | Total Landsman | % 1971 | Large vessel | Landsman | | | | Total Landsman | % 1972 | | | | |
|-----|--------------|-----|----------|----|----|----|----------------|--------|--------------|----------|----|----|----|----------------|--------|--------------|----------|----|---|---|----------------|--------|---|---|---|---|
| | A | B | A | B | C | D | | | | A | B | C | D | | | | A | B | C | D | | | A | B | C | D |
| | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | 105 | 134 | 239 | 41 | 31 | 5 | 21 | 98 | 18.17 | 239 | 3 | 4 | 17 | 39 | 63 | 31.59 | 12 | 11 | 1 | 0 | 12 | 6.21 | | | | |
| 2 | 98 | 112 | 210 | 41 | 69 | 12 | 60 | 182 | 17.46 | 66 | 18 | 15 | 29 | 27 | 89 | 10.72 | 9 | 19 | 6 | 3 | 28 | 8.10 | | | | |
| 3 | 11 | 34 | 45 | 15 | 89 | 32 | 67 | 203 | 6.29 | 53 | 35 | 51 | 45 | 33 | 164 | 11.19 | 9 | 17 | 2 | 1 | 20 | 14.27 | | | | |
| 4 | 17 | 29 | 46 | 25 | 45 | 55 | 63 | 188 | 6.13 | 19 | 34 | 30 | 17 | 14 | 95 | 5.02 | 24 | 26 | 3 | 3 | 32 | 9.88 | | | | |
| 5 | 18 | 15 | 33 | 23 | 40 | 77 | 68 | 208 | 5.53 | 18 | 45 | 32 | 31 | 5 | 113 | 5.37 | 9 | 10 | 1 | 6 | 17 | 10.07 | | | | |
| 6 | 12 | 18 | 30 | 19 | 16 | 38 | 44 | 117 | 3.91 | 11 | 30 | 9 | 9 | 21 | 69 | 3.30 | 10 | 15 | 0 | 3 | 18 | 6.54 | | | | |
| 7 | 15 | 26 | 41 | 14 | 8 | 35 | 24 | 81 | 4.12 | 10 | 16 | 9 | 6 | 18 | 49 | 2.61 | 13 | 5 | 0 | 4 | 9 | 5.15 | | | | |
| 8 | 8 | 15 | 23 | 43 | 12 | 64 | 22 | 141 | 3.79 | 8 | 23 | 4 | 6 | 15 | 48 | 2.34 | 9 | 8 | 0 | 3 | 11 | 4.72 | | | | |
| 9 | 13 | 22 | 35 | 49 | 8 | 66 | 25 | 148 | 4.74 | 17 | 21 | 7 | 3 | 13 | 44 | 3.35 | 6 | 5 | 1 | 0 | 6 | 4.14 | | | | |
| 10 | 18 | 13 | 31 | 49 | 8 | 82 | 24 | 153 | 4.54 | 16 | 30 | 3 | 4 | 31 | 58 | 3.61 | 6 | 6 | 0 | 1 | 7 | 5.29 | | | | |
| 11 | 13 | 7 | 20 | 24 | 4 | 61 | 13 | 102 | 2.97 | 10 | 9 | 2 | 10 | 20 | 41 | 2.39 | 4 | 13 | 0 | 2 | 15 | 3.72 | | | | |
| 12 | 20 | 6 | 26 | 28 | 4 | 42 | 8 | 82 | 3.09 | 12 | 6 | 6 | 2 | 26 | 40 | 2.61 | 5 | 7 | 0 | 6 | 13 | 3.72 | | | | |
| 13 | 12 | 11 | 23 | 26 | 1 | 37 | 8 | 72 | 2.72 | 8 | 8 | 6 | 4 | 22 | 40 | 2.11 | 2 | 5 | 0 | 2 | 7 | 3.46 | | | | |
| 14 | 6 | 10 | 16 | 19 | 0 | 30 | 5 | 54 | 1.95 | 4 | 9 | 3 | 0 | 13 | 25 | 1.20 | 2 | 15 | 0 | 0 | 15 | 2.23 | | | | |
| 15 | 11 | 13 | 24 | 29 | 2 | 32 | 10 | 73 | 2.81 | 6 | 4 | 3 | 0 | 18 | 25 | 1.45 | 6 | 6 | 0 | 2 | 8 | 2.57 | | | | |
| 16 | 13 | 4 | 17 | 30 | 1 | 27 | 2 | 60 | 2.12 | 6 | 3 | 5 | 0 | 11 | 19 | 1.28 | 4 | 3 | 1 | 0 | 4 | 1.91 | | | | |
| 17 | 7 | 8 | 15 | 20 | 0 | 30 | 9 | 59 | 1.96 | 6 | 5 | 3 | 1 | 8 | 17 | 1.22 | 3 | 8 | 0 | 0 | 8 | 1.66 | | | | |
| 18 | 7 | 10 | 17 | 12 | 1 | 21 | 6 | 40 | 1.81 | 9 | 4 | 1 | 0 | 10 | 15 | 1.54 | 2 | 7 | 0 | 3 | 10 | 1.41 | | | | |
| 19 | 7 | 7 | 14 | 18 | 1 | 17 | 6 | 42 | 1.63 | 6 | 3 | 2 | 0 | 7 | 12 | 1.08 | 0 | 5 | 0 | 0 | 5 | 0.99 | | | | |
| 20 | 6 | 6 | 12 | 22 | 3 | 16 | 6 | 47 | 1.56 | 8 | 2 | 0 | 1 | 11 | 14 | 1.39 | 5 | 13 | 0 | 1 | 14 | 1.58 | | | | |
| 21 | 4 | 4 | 8 | 10 | 1 | 10 | 1 | 22 | 0.90 | 5 | 0 | 0 | 2 | 7 | 9 | 0.87 | 1 | 7 | 1 | 1 | 9 | 0.83 | | | | |
| 22 | 5 | 2 | 7 | 5 | 0 | 9 | 1 | 15 | 0.72 | 9 | 0 | 0 | 0 | 4 | 4 | 1.23 | 3 | 3 | 0 | 2 | 5 | 0.59 | | | | |
| 23 | 3 | 0 | 3 | 7 | 0 | 4 | 3 | 14 | 0.43 | 7 | 0 | 0 | 0 | 4 | 4 | 0.99 | 1 | 2 | 0 | 2 | 2 | 0.41 | | | | |
| 24 | 3 | 1 | 4 | 5 | 0 | 5 | 1 | 11 | 0.45 | 4 | 1 | 0 | 0 | 1 | 2 | 0.56 | 2 | 3 | 0 | 0 | 3 | 0.34 | | | | |
| 25 | 0 | 1 | 1 | 3 | 0 | 6 | 0 | 9 | 0.21 | 7 | 1 | 0 | 0 | 2 | 3 | 0.96 | 0 | 0 | 0 | 0 | 0 | 0.25 | | | | |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | | Total | | Landsman | | % 1973 | | Large vessel | | Landsman | | | | Total | | % 1974 | | Large vessel | | Landsman | | | | Total | | % 1975 | | | | | | | | |
|-----|--------------|---|--------------|----|----------|-----|--------|----|--------------|-----|----------|-------|-----|----|-------|-----|--------|----|--------------|-------|----------|-------|---|---|-------|---|--------|---|-------|---|---|---|---|---|---|
| | A | B | Large vessel | A | A | B | C | D | Total | A | A | B | C | D | Total | A | A | B | C | D | E | Total | A | A | B | C | D | E | Total | A | A | B | C | D | E |
| 1 | 171 | 4 | 175 | 6 | 6.52 | 360 | 0 | 8 | 46 | 106 | 160 | 26.66 | 464 | 20 | 0 | 157 | 146 | 55 | 378 | 28.80 | | | | | | | | | | | | | | | |
| 2 | 232 | 5 | 237 | 14 | 9.61 | 227 | 0 | 18 | 33 | 22 | 73 | 15.62 | 223 | 55 | 1 | 123 | 30 | 47 | 256 | 17.10 | | | | | | | | | | | | | | | |
| 3 | 188 | 0 | 188 | 16 | 8.40 | 86 | 2 | 7 | 11 | 7 | 27 | 5.89 | 102 | 56 | 12 | 61 | 13 | 20 | 162 | 9.79 | | | | | | | | | | | | | | | |
| 4 | 173 | 5 | 178 | 9 | 6.98 | 65 | 5 | 10 | 10 | 5 | 30 | 4.86 | 67 | 19 | 26 | 40 | 1 | 10 | 96 | 5.98 | | | | | | | | | | | | | | | |
| 5 | 293 | 6 | 299 | 11 | 11.26 | 67 | 17 | 12 | 5 | 6 | 40 | 5.40 | 57 | 8 | 62 | 29 | 4 | 4 | 107 | 6.20 | | | | | | | | | | | | | | | |
| 6 | 81 | 0 | 81 | 11 | 4.16 | 88 | 61 | 19 | 7 | 4 | 91 | 8.73 | 57 | 4 | 57 | 25 | 2 | 5 | 93 | 5.58 | | | | | | | | | | | | | | | |
| 7 | 67 | 1 | 68 | 16 | 4.34 | 21 | 20 | 13 | 3 | 1 | 37 | 2.73 | 28 | 5 | 71 | 22 | 1 | 9 | 108 | 5.48 | | | | | | | | | | | | | | | |
| 8 | 79 | 6 | 85 | 15 | 4.61 | 21 | 30 | 18 | 3 | 0 | 51 | 3.33 | 23 | 4 | 50 | 11 | 0 | 2 | 67 | 3.55 | | | | | | | | | | | | | | | |
| 9 | 42 | 0 | 42 | 21 | 4.16 | 36 | 19 | 17 | 1 | 0 | 37 | 3.56 | 29 | 3 | 21 | 7 | 0 | 2 | 33 | 2.21 | | | | | | | | | | | | | | | |
| 10 | 37 | 4 | 41 | 17 | 3.47 | 21 | 15 | 20 | 0 | 0 | 35 | 2.65 | 17 | 0 | 19 | 18 | 0 | 10 | 47 | 2.07 | | | | | | | | | | | | | | | |
| 11 | 45 | 3 | 48 | 24 | 4.65 | 16 | 8 | 18 | 0 | 1 | 27 | 2.03 | 28 | 0 | 14 | 14 | 0 | 1 | 29 | 2.01 | | | | | | | | | | | | | | | |
| 12 | 30 | 0 | 30 | 27 | 4.54 | 24 | 11 | 14 | 2 | 0 | 27 | 2.47 | 17 | 2 | 14 | 11 | 0 | 4 | 31 | 1.81 | | | | | | | | | | | | | | | |
| 13 | 36 | 0 | 36 | 27 | 4.09 | 23 | 6 | 17 | 1 | 0 | 24 | 2.29 | 20 | 0 | 10 | 20 | 1 | 1 | 32 | 1.93 | | | | | | | | | | | | | | | |
| 14 | 24 | 0 | 24 | 22 | 3.56 | 19 | 2 | 17 | 1 | 0 | 20 | 1.90 | 15 | 0 | 9 | 18 | 0 | 0 | 27 | 1.58 | | | | | | | | | | | | | | | |
| 15 | 36 | 1 | 37 | 18 | 3.56 | 30 | 2 | 16 | 0 | 0 | 18 | 1.86 | 15 | 0 | 5 | 10 | 0 | 0 | 15 | 1.05 | | | | | | | | | | | | | | | |
| 16 | 32 | 1 | 33 | 15 | 3.04 | 16 | 2 | 21 | 0 | 1 | 24 | 1.90 | 13 | 0 | 7 | 7 | 0 | 2 | 16 | 1.04 | | | | | | | | | | | | | | | |
| 17 | 25 | 1 | 26 | 11 | 2.28 | 16 | 4 | 13 | 1 | 0 | 18 | 1.65 | 13 | 0 | 1 | 7 | 0 | 1 | 9 | 0.74 | | | | | | | | | | | | | | | |
| 18 | 23 | 1 | 24 | 10 | 2.08 | 14 | 1 | 10 | 0 | 0 | 11 | 1.24 | 10 | 0 | 7 | 9 | 0 | 0 | 16 | 0.97 | | | | | | | | | | | | | | | |
| 19 | 29 | 1 | 30 | 6 | 1.76 | 12 | 2 | 7 | 0 | 0 | 9 | 1.04 | 8 | 0 | 2 | 4 | 0 | 0 | 6 | 0.47 | | | | | | | | | | | | | | | |
| 20 | 33 | 0 | 33 | 6 | 1.89 | 10 | 1 | 6 | 0 | 0 | 7 | 0.85 | 6 | 0 | 0 | 4 | 0 | 1 | 5 | 0.38 | | | | | | | | | | | | | | | |
| 21 | 18 | 0 | 18 | 3 | 0.99 | 8 | 0 | 8 | 0 | 0 | 8 | 0.78 | 5 | 0 | 0 | 3 | 0 | 0 | 3 | 0.26 | | | | | | | | | | | | | | | |
| 22 | 18 | 0 | 18 | 13 | 0.23 | 9 | 0 | 7 | 0 | 0 | 7 | 0.79 | 7 | 0 | 1 | 2 | 0 | 0 | 3 | 0.32 | | | | | | | | | | | | | | | |
| 23 | 20 | 0 | 20 | 0 | 0.67 | 7 | 0 | 4 | 0 | 0 | 4 | 0.56 | 5 | 0 | 0 | 0 | 0 | 1 | 1 | 0.18 | | | | | | | | | | | | | | | |
| 24 | 15 | 0 | 15 | 0 | 0.50 | 11 | 0 | 0 | 0 | 0 | 0 | 0.61 | 6 | 0 | 1 | 2 | 0 | 0 | 3 | 0.29 | | | | | | | | | | | | | | | |
| 25 | 17 | 0 | 17 | 0 | 0.56 | 9 | 1 | 1 | 0 | 0 | 2 | 0.58 | 5 | 0 | 0 | 2 | 0 | 0 | 2 | 0.22 | | | | | | | | | | | | | | | |

Table 1. (cont'd) Catch at age of the northwest Atlantic harp seal from different landsmen and large vessel fisheries. Individual samples are shown from 1961 on. Annual percentage compositions have been weighted in relation to the ratio of landsmen to larger vessel catch.

| Age | Large vessel | Landsman | | | | | | | Total Landsman | Total \$ 1976 |
|-----|--------------|----------|----|----|----|-----|--|-----|----------------|---------------|
| | | A | B | C | D | E | | | | |
| 1 | 292 | 0 | 7 | 88 | 75 | 113 | | 283 | 33.20 | |
| 2 | 112 | 0 | 18 | 57 | 34 | 58 | | 167 | 18.96 | |
| 3 | 60 | 1 | 26 | 45 | 20 | 25 | | 117 | 12.08 | |
| 4 | 35 | 13 | 9 | 37 | 11 | 17 | | 87 | 8.73 | |
| 5 | 24 | 16 | 4 | 22 | 2 | 2 | | 46 | 4.76 | |
| 6 | 13 | 15 | 3 | 20 | 1 | 0 | | 39 | 3.84 | |
| 7 | 18 | 19 | 1 | 10 | 1 | 0 | | 31 | 3.26 | |
| 8 | 13 | 13 | 0 | 16 | 2 | 0 | | 31 | 3.12 | |
| 9 | 11 | 9 | 1 | 3 | 0 | 0 | | 13 | 1.46 | |
| 10 | 6 | 14 | 1 | 9 | 0 | 0 | | 24 | 2.30 | |
| 11 | 6 | 8 | 0 | 6 | 2 | 0 | | 16 | 1.59 | |
| 12 | 5 | 4 | 0 | 5 | 2 | 0 | | 11 | 1.12 | |
| 13 | 8 | 5 | 0 | 3 | 0 | 0 | | 8 | 0.93 | |
| 14 | 5 | 4 | 0 | 4 | 1 | 0 | | 9 | 0.94 | |
| 15 | 5 | 3 | 0 | 13 | 0 | 0 | | 16 | 1.57 | |
| 16 | 5 | 1 | 0 | 4 | 0 | 0 | | 5 | 0.58 | |
| 17 | 5 | 1 | 0 | 5 | 0 | 0 | | 6 | 0.67 | |
| 18 | 3 | 0 | 0 | 4 | 0 | 0 | | 4 | 0.44 | |
| 19 | 2 | 0 | 0 | 4 | 0 | 0 | | 4 | 0.41 | |
| 20 | 1 | 0 | 0 | 5 | 0 | 0 | | 5 | 0.47 | |
| 21 | 1 | 0 | 0 | 1 | 0 | 0 | | 1 | 0.12 | |
| 22 | 2 | 0 | 0 | 1 | 0 | 0 | | 1 | 0.14 | |
| 23 | 1 | 0 | 0 | 1 | 0 | 0 | | 1 | 0.16 | |
| 24 | 0 | 0 | 0 | 1 | 0 | 0 | | 1 | 0.09 | |
| 25 | 0 | 0 | 0 | 1 | 0 | 0 | | 1 | 0.09 | |

Table 2. The total catch of 1+ northwest Atlantic harp seals by landsmen and large vessels from 1952 to 1975.

| YEAR | LANDSMEN | LARGE VESSEL | % LANDSMEN | % LARGE VESSEL |
|------|----------|--------------|------------|----------------|
| 1952 | 10667 | 98378 | 9.8 | 90.2 |
| 53 | 8100 | 66811 | 10.8 | 89.2 |
| 54 | 5443 | 83939 | 6.1 | 93.9 |
| 1955 | 5401 | 75671 | 6.7 | 93.3 |
| 56 | 5428 | 42585 | 11.3 | 88.7 |
| 57 | 3605 | 76437 | 4.5 | 95.5 |
| 58 | 19563 | 137227 | 12.5 | 87.5 |
| 59 | 3998 | 77304 | 4.9 | 95.1 |
| 1960 | 6648 | 114534 | 5.5 | 94.5 |
| 61 | 5877 | 13170 | 30.9 | 69.1 |
| 62 | 13388 | 99513 | 11.9 | 88.1 |
| 63 | 14529 | 57094 | 20.3 | 79.7 |
| 64 | 14933 | 60348 | 19.8 | 80.2 |
| 1965 | 17738 | 33757 | 34.4 | 65.6 |
| 66 | 12647 | 59364 | 17.6 | 82.4 |
| 67 | 15245 | 41361 | 26.9 | 73.1 |
| 68 | 5910 | 30328 | 16.3 | 83.7 |
| 69 | 10532 | 44940 | 19.0 | 81.0 |
| 1970 | 13839 | 26225 | 34.5 | 65.5 |
| 71 | 6044 | 14343 | 29.6 | 70.4 |
| 72 | 11427 | 1646 | 87.4 | 12.6 |
| 73 | 10416 | 15081 | 40.9 | 59.1 |
| 74 | 10982 | 21828 | 33.5 | 66.5 |
| 1975 | 22733 | 10992 | 67.4 | 32.6 |

Table 3. Catch at age data of the northwest Atlantic harp seals resulting from the weighting of samples and application to the total catch.

| Age | 1952 | 1953 | 1954 | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 198063 | 197975 | 184491 | 260020 | 346846 | 171909 | 149350 | 243255 | 164158 | 174762 | 211285 | 285994 |
| 1 | 4679 | 20685 | 35353 | 26005 | 14664 | 25488 | 44871 | 25180 | 37848 | 6586 | 27588 | 6780 |
| 2 | 11529 | 6527 | 14119 | 12499 | 7175 | 11796 | 20701 | 11778 | 18220 | 2597 | 34154 | 8785 |
| 3 | 7589 | 5888 | 4257 | 7526 | 4215 | 6631 | 12830 | 6675 | 10108 | 2566 | 9762 | 8044 |
| 4 | 5983 | 3913 | 6444 | 5467 | 3355 | 5280 | 11387 | 5194 | 9486 | 3155 | 9549 | 4980 |
| 5 | 6875 | 3988 | 3590 | 4738 | 2660 | 4248 | 9163 | 4360 | 6478 | 1067 | 7108 | 4215 |
| 6 | 12029 | 3207 | 4207 | 3889 | 2310 | 3647 | 8247 | 3838 | 5783 | 1335 | 2932 | 4572 |
| 7 | 8451 | 2843 | 3879 | 3367 | 2108 | 3256 | 6812 | 3365 | 4877 | 1326 | 2978 | 4308 |
| 8 | 8302 | 2790 | 3351 | 3171 | 2045 | 2908 | 6268 | 3021 | 4200 | 811 | 3081 | 4349 |
| 9 | 5889 | 2732 | 2246 | 2588 | 1804 | 2618 | 5347 | 2731 | 3737 | 771 | 2963 | 3993 |
| 10 | 6711 | 2263 | 3271 | 2414 | 1588 | 2493 | 5231 | 2485 | 3506 | 1192 | 1350 | 4582 |
| 11 | 5889 | 2741 | 1350 | 2159 | 1378 | 2285 | 4754 | 2168 | 3145 | 712 | 1462 | 4440 |
| 12 | 1740 | 1664 | 2841 | 2031 | 1349 | 2088 | 4129 | 1961 | 2896 | 409 | 2075 | 3272 |
| 13 | 1532 | 1267 | 2806 | 1763 | 1169 | 1877 | 3792 | 1792 | 2627 | 411 | 1160 | 3440 |
| 14 | 2325 | 996 | 1430 | 1659 | 1112 | 1809 | 3419 | 1741 | 2488 | 403 | 1550 | 3562 |
| 15 | 4208 | 1909 | 1882 | 1567 | 964 | 1670 | 3141 | 1595 | 2264 | 204 | 2319 | 2828 |
| 16 | 1443 | 1901 | 2415 | 1485 | 1006 | 1541 | 2774 | 1479 | 2172 | 325 | 810 | 2240 |
| 17 | 2355 | 1395 | 1092 | 1311 | 867 | 1367 | 2566 | 1319 | 1912 | 248 | 1829 | 2355 |
| 18 | 1844 | 879 | 391 | 1161 | 865 | 1205 | 2188 | 1128 | 1673 | 130 | 914 | 1522 |
| 19 | 962 | 640 | 1243 | 1068 | 734 | 1112 | 1916 | 1075 | 1538 | 157 | 652 | 1242 |
| 20 | 4971 | 2786 | 746 | 859 | 624 | 964 | 1741 | 854 | 1273 | 193 | 1470 | 1306 |
| 21 | 1383 | 1630 | 356 | 674 | 450 | 696 | 1150 | 617 | 923 | 53 | 183 | 1107 |
| 22 | 60 | 909 | 338 | 592 | 392 | 615 | 1028 | 548 | 831 | 105 | 715 | 865 |
| 23 | 481 | 649 | 773 | 418 | 321 | 417 | 842 | 427 | 616 | 78 | 199 | 717 |
| 24 | 1344 | 468 | 151 | 348 | 263 | 371 | 702 | 343 | 513 | 65 | 183 | 566 |
| 25 | 481 | 254 | 320 | 47 | 54 | 83 | 157 | 62 | 69 | 103 | 215 | 267 |

Table 3. (cont'd) Catch at age data of the northwest Atlantic harp seal resulting from the weighting of samples and application to the total catch.

| Age | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 270952 | 187284 | 225250 | 279858 | 159971 | 236532 | 220520 | 213349 | 119658 | 102744 | 118036 | 140629 |
| 1 | 2502 | 7413 | 11558 | 10877 | 5980 | 20985 | 7844 | 7315 | 1990 | 1949 | 9604 | 9629 |
| 2 | 4418 | 3693 | 10383 | 5100 | 4521 | 2993 | 7535 | 2483 | 1289 | 2874 | 5629 | 5719 |
| 3 | 6364 | 4898 | 5183 | 1959 | 3069 | 3088 | 2714 | 2591 | 2272 | 2511 | 2123 | 3275 |
| 4 | 7429 | 6408 | 5414 | 2090 | 1823 | 2512 | 2644 | 1163 | 1573 | 2087 | 1752 | 1999 |
| 5 | 5364 | 7889 | 5880 | 3587 | 1682 | 2966 | 2387 | 1250 | 1604 | 3367 | 1945 | 2072 |
| 6 | 7702 | 7349 | 6086 | 4760 | 1711 | 2125 | 1687 | 764 | 1041 | 1243 | 3144 | 1867 |
| 7 | 4117 | 3121 | 5373 | 4341 | 2437 | 2479 | 1776 | 606 | 820 | 1299 | 985 | 1832 |
| 8 | 3419 | 1826 | 3543 | 3059 | 2498 | 2983 | 1637 | 542 | 752 | 1380 | 1200 | 1186 |
| 9 | 3189 | 1081 | 2018 | 2227 | 1705 | 2325 | 2045 | 776 | 659 | 1245 | 1283 | 740 |
| 10 | 4751 | 1524 | 1928 | 1830 | 1757 | 1831 | 1958 | 837 | 842 | 1038 | 954 | 693 |
| 11 | 2597 | 655 | 2522 | 2282 | 1310 | 1666 | 1286 | 554 | 592 | 1392 | 732 | 672 |
| 12 | 2544 | 2088 | 1756 | 1570 | 1031 | 1153 | 1332 | 606 | 593 | 1360 | 891 | 605 |
| 13 | 2023 | 660 | 1639 | 1316 | 976 | 1295 | 1175 | 490 | 552 | 1224 | 825 | 646 |
| 14 | 1988 | 1553 | 1825 | 1852 | 1233 | 1333 | 844 | 277 | 355 | 1064 | 685 | 529 |
| 15 | 2799 | 1415 | 1548 | 1810 | 1082 | 1393 | 1211 | 335 | 410 | 1067 | 674 | 353 |
| 16 | 2314 | 794 | 1580 | 1289 | 1037 | 991 | 914 | 297 | 304 | 909 | 686 | 350 |
| 17 | 2887 | 512 | 1104 | 1964 | 903 | 1248 | 847 | 284 | 264 | 682 | 594 | 247 |
| 18 | 4290 | 846 | 1619 | 1592 | 1237 | 1039 | 780 | 357 | 224 | 623 | 447 | 323 |
| 19 | 2331 | 961 | 1035 | 1608 | 1073 | 1042 | 703 | 251 | 158 | 526 | 377 | 159 |
| 20 | 142 | 340 | 1094 | 1143 | 742 | 927 | 676 | 322 | 252 | 566 | 306 | 127 |
| 21 | 2189 | 593 | 780 | 684 | 485 | 680 | 389 | 203 | 132 | 298 | 282 | 88 |
| 22 | 1130 | 282 | 349 | 491 | 684 | 554 | 312 | 286 | 94 | 690 | 287 | 106 |
| 23 | 1130 | 58 | 687 | 552 | 379 | 491 | 185 | 229 | 67 | 201 | 201 | 59 |
| 24 | 1616 | 87 | 373 | 400 | 267 | 267 | 195 | 129 | 54 | 151 | 219 | 97 |
| 25 | 628 | 87 | 256 | 342 | 139 | 310 | 91 | 222 | 40 | 171 | 210 | 74 |

Table 4. Hunting mortality levels of the northwest Atlantic harp seal calculated from cohort analysis assuming $M = 0.114$.

| Age | 1952 | 1953 | 1954 | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0 | 0.644 | 0.585 | 0.634 | 0.844 | 0.989 | 0.499 | 0.524 | 0.733 | 0.557 | 0.570 | 0.759 | 1.101 |
| 1 | 0.024 | 0.112 | 0.174 | 0.151 | 0.088 | 0.150 | 0.210 | 0.140 | 0.209 | 0.034 | 0.146 | 0.042 |
| 2 | 0.062 | 0.039 | 0.095 | 0.078 | 0.052 | 0.087 | 0.159 | 0.072 | 0.130 | 0.018 | 0.218 | 0.058 |
| 3 | 0.032 | 0.037 | 0.029 | 0.062 | 0.032 | 0.057 | 0.116 | 0.065 | 0.074 | 0.022 | 0.075 | 0.067 |
| 4 | 0.032 | 0.019 | 0.047 | 0.044 | 0.033 | 0.046 | 0.119 | 0.058 | 0.112 | 0.027 | 0.096 | 0.046 |
| 5 | 0.060 | 0.024 | 0.020 | 0.041 | 0.025 | 0.048 | 0.095 | 0.056 | 0.087 | 0.015 | 0.071 | 0.051 |
| 6 | 0.066 | 0.033 | 0.030 | 0.025 | 0.023 | 0.039 | 0.112 | 0.048 | 0.089 | 0.021 | 0.047 | 0.055 |
| 7 | 0.084 | 0.019 | 0.047 | 0.027 | 0.016 | 0.038 | 0.086 | 0.056 | 0.072 | 0.024 | 0.053 | 0.082 |
| 8 | 0.055 | 0.033 | 0.025 | 0.045 | 0.019 | 0.024 | 0.086 | 0.046 | 0.084 | 0.014 | 0.066 | 0.092 |
| 9 | 0.066 | 0.022 | 0.031 | 0.022 | 0.030 | 0.028 | 0.051 | 0.045 | 0.067 | 0.018 | 0.060 | 0.103 |
| 10 | 0.049 | 0.030 | 0.029 | 0.038 | 0.016 | 0.048 | 0.065 | 0.028 | 0.068 | 0.025 | 0.036 | 0.112 |
| 11 | 0.063 | 0.023 | 0.021 | 0.022 | 0.025 | 0.025 | 0.110 | 0.032 | 0.041 | 0.016 | 0.035 | 0.145 |
| 12 | 0.035 | 0.021 | 0.028 | 0.036 | 0.016 | 0.044 | 0.053 | 0.055 | 0.049 | 0.006 | 0.052 | 0.091 |
| 13 | 0.014 | 0.030 | 0.041 | 0.020 | 0.024 | 0.025 | 0.096 | 0.027 | 0.089 | 0.008 | 0.019 | 0.103 |
| 14 | 0.038 | 0.010 | 0.039 | 0.028 | 0.014 | 0.042 | 0.053 | 0.054 | 0.043 | 0.016 | 0.033 | 0.068 |
| 15 | 0.093 | 0.036 | 0.022 | 0.050 | 0.019 | 0.024 | 0.087 | 0.029 | 0.083 | 0.004 | 0.109 | 0.070 |
| 16 | 0.058 | 0.051 | 0.054 | 0.019 | 0.038 | 0.034 | 0.047 | 0.050 | 0.046 | 0.014 | 0.018 | 0.132 |
| 17 | 0.133 | 0.066 | 0.034 | 0.034 | 0.013 | 0.060 | 0.067 | 0.026 | 0.076 | 0.006 | 0.088 | 0.059 |
| 18 | 0.140 | 0.061 | 0.022 | 0.042 | 0.026 | 0.020 | 0.117 | 0.035 | 0.038 | 0.006 | 0.024 | 0.090 |
| 19 | 0.045 | 0.061 | 0.105 | 0.070 | 0.031 | 0.039 | 0.037 | 0.071 | 0.056 | 0.004 | 0.032 | 0.037 |
| 20 | 0.383 | 0.159 | 0.085 | 0.090 | 0.049 | 0.048 | 0.071 | 0.019 | 0.102 | 0.008 | 0.043 | 0.075 |
| 21 | 0.186 | 0.188 | 0.025 | 0.094 | 0.057 | 0.064 | 0.067 | 0.030 | 0.024 | 0.005 | 0.008 | 0.038 |
| 22 | 0.014 | 0.163 | 0.050 | 0.049 | 0.066 | 0.094 | 0.115 | 0.038 | 0.047 | 0.003 | 0.077 | 0.044 |
| 23 | 0.023 | 0.188 | 0.185 | 0.073 | 0.031 | 0.085 | 0.163 | 0.059 | 0.050 | 0.005 | 0.005 | 0.094 |
| 24 | 0.090 | 0.026 | 0.056 | 0.108 | 0.055 | 0.041 | 0.183 | 0.085 | 0.085 | 0.006 | 0.013 | 0.016 |
| 25 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 |

Table 4. (cont'd) Hunting mortality levels of the northwest Atlantic harp seal calculated from cohort analysis assuming $M = 0.114$.

| Age | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0 | 1.038 | 0.861 | 1.019 | 1.039 | 0.528 | 0.826 | 0.809 | 0.900 | 0.448 | 0.426 | 0.428 |
| 1 | 0.020 | 0.058 | 0.099 | 0.101 | 0.045 | 0.108 | 0.049 | 0.048 | 0.008 | 0.011 | 0.058 |
| 2 | 0.032 | 0.034 | 0.098 | 0.053 | 0.051 | 0.026 | 0.047 | 0.018 | 0.010 | 0.026 | 0.035 |
| 3 | 0.050 | 0.041 | 0.056 | 0.022 | 0.038 | 0.041 | 0.028 | 0.019 | 0.019 | 0.022 | 0.022 |
| 4 | 0.074 | 0.059 | 0.053 | 0.027 | 0.024 | 0.036 | 0.041 | 0.014 | 0.013 | 0.020 | 0.018 |
| 5 | 0.059 | 0.096 | 0.065 | 0.042 | 0.025 | 0.045 | 0.040 | 0.023 | 0.021 | 0.032 | 0.021 |
| 6 | 0.113 | 0.097 | 0.091 | 0.063 | 0.023 | 0.036 | 0.030 | 0.015 | 0.022 | 0.019 | 0.035 |
| 7 | 0.058 | 0.056 | 0.087 | 0.079 | 0.038 | 0.038 | 0.035 | 0.012 | 0.018 | 0.031 | 0.017 |
| 8 | 0.079 | 0.030 | 0.076 | 0.060 | 0.055 | 0.054 | 0.030 | 0.012 | 0.017 | 0.035 | 0.033 |
| 9 | 0.083 | 0.030 | 0.039 | 0.057 | 0.040 | 0.060 | 0.044 | 0.016 | 0.017 | 0.033 | 0.037 |
| 10 | 0.156 | 0.048 | 0.062 | 0.041 | 0.054 | 0.050 | 0.061 | 0.021 | 0.020 | 0.030 | 0.029 |
| 11 | 0.078 | 0.027 | 0.094 | 0.089 | 0.034 | 0.060 | 0.041 | 0.020 | 0.017 | 0.038 | 0.025 |
| 12 | 0.106 | 0.076 | 0.084 | 0.071 | 0.048 | 0.035 | 0.057 | 0.023 | 0.025 | 0.045 | 0.028 |
| 13 | 0.069 | 0.033 | 0.072 | 0.077 | 0.053 | 0.072 | 0.041 | 0.025 | 0.024 | 0.059 | 0.032 |
| 14 | 0.073 | 0.063 | 0.109 | 0.099 | 0.087 | 0.087 | 0.056 | 0.012 | 0.021 | 0.052 | 0.039 |
| 15 | 0.064 | 0.062 | 0.075 | 0.137 | 0.071 | 0.123 | 0.097 | 0.026 | 0.019 | 0.072 | 0.039 |
| 16 | 0.069 | 0.021 | 0.984 | 0.076 | 0.099 | 0.078 | 0.101 | 0.029 | 0.027 | 0.048 | 0.055 |
| 17 | 0.228 | 0.018 | 0.034 | 0.129 | 0.064 | 0.151 | 0.081 | 0.038 | 0.029 | 0.071 | 0.037 |
| 18 | 0.132 | 0.088 | 0.066 | 0.057 | 0.102 | 0.089 | 0.122 | 0.041 | 0.035 | 0.081 | 0.056 |
| 19 | 0.175 | 0.036 | 0.135 | 0.079 | 0.046 | 0.107 | 0.073 | 0.048 | 0.021 | 0.097 | 0.059 |
| 20 | 0.005 | 0.032 | 0.048 | 0.197 | 0.044 | 0.046 | 0.086 | 0.040 | 0.057 | 0.088 | 0.069 |
| 21 | 0.158 | 0.023 | 0.087 | 0.035 | 0.109 | 0.047 | 0.023 | 0.031 | 0.019 | 0.080 | 0.053 |
| 22 | 0.045 | 0.025 | 0.016 | 0.066 | 0.041 | 0.160 | 0.025 | 0.019 | 0.017 | 0.117 | 0.095 |
| 23 | 0.067 | 0.003 | 0.072 | 0.028 | 0.061 | 0.034 | 0.067 | 0.021 | 0.005 | 0.040 | 0.042 |
| 24 | 0.284 | 0.006 | 0.020 | 0.050 | 0.016 | 0.051 | 0.016 | 0.056 | 0.006 | 0.013 | 0.052 |
| 25 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 | 0.020 |

Table 5. Numbers at age of the northwest Atlantic harp seal estimated by cohort analysis from catch data (Table 3) assuming $M = 0.114$.

| Age | 1952 | 1953 | 1954 | 1955 | 1956 | 1957 | 1958 | 1959 | 1960 | 1961 | 1962 | 1963 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 441614 | 473555 | 416283 | 482960 | 584822 | 463624 | 387829 | 495879 | 407345 | 426122 | 420630 | 453722 |
| 1 | 211501 | 206945 | 235528 | 197163 | 185312 | 194184 | 251288 | 204969 | 212675 | 214060 | 215132 | 175732 |
| 2 | 205981 | 184294 | 165109 | 176757 | 151356 | 151495 | 149186 | 181829 | 159100 | 162175 | 184776 | 165893 |
| 3 | 259377 | 172898 | 158272 | 133983 | 145907 | 128271 | 124030 | 113559 | 151113 | 127367 | 142249 | 132606 |
| 4 | 205951 | 224263 | 148708 | 137199 | 112439 | 126205 | 108187 | 98548 | 95018 | 127365 | 111220 | 117787 |
| 5 | 125093 | 178110 | 196404 | 126599 | 117252 | 97155 | 107620 | 85775 | 83024 | 77614 | 110663 | 90217 |
| 6 | 201628 | 105121 | 155153 | 171852 | 108484 | 102107 | 82675 | 87370 | 72415 | 70886 | 68244 | 92025 |
| 7 | 112067 | 168542 | 90766 | 134462 | 149663 | 94613 | 87661 | 65978 | 74331 | 59444 | 61988 | 58122 |
| 8 | 164385 | 92010 | 147697 | 77322 | 116795 | 131547 | 81344 | 71782 | 55690 | 62201 | 51787 | 52496 |
| 9 | 98334 | 138832 | 79461 | 128619 | 65996 | 102279 | 114627 | 66659 | 61194 | 46345 | 54733 | 43297 |
| 10 | 151167 | 82177 | 121293 | 68778 | 112317 | 57182 | 88787 | 97226 | 56898 | 53117 | 40623 | 46037 |
| 11 | 102492 | 128541 | 71185 | 105135 | 59088 | 98715 | 48666 | 74280 | 84403 | 50227 | 46269 | 34971 |
| 12 | 53662 | 85887 | 112103 | 62241 | 91768 | 51420 | 85921 | 38932 | 64229 | 74773 | 44143 | 39903 |
| 13 | 123154 | 46237 | 75061 | 97341 | 53616 | 80607 | 43907 | 72764 | 32885 | 58040 | 66330 | 37427 |
| 14 | 66875 | 108438 | 40059 | 64324 | 85188 | 46735 | 70149 | 35595 | 63231 | 27271 | 51399 | 58088 |
| 15 | 50322 | 57474 | 95814 | 34392 | 55826 | 74959 | 39991 | 59362 | 30115 | 56353 | 23952 | 44397 |
| 16 | 27505 | 40925 | 49478 | 83713 | 29206 | 48901 | 65306 | 32716 | 51459 | 26229 | 50089 | 19181 |
| 17 | 20122 | 23178 | 34720 | 41866 | 73291 | 25109 | 42177 | 55649 | 27794 | 47867 | 23096 | 43927 |
| 18 | 14974 | 15729 | 19363 | 29948 | 36117 | 64575 | 21113 | 35209 | 48408 | 25001 | 42476 | 18880 |
| 19 | 23533 | 11619 | 13205 | 16908 | 25625 | 31409 | 56480 | 16771 | 30350 | 42072 | 22185 | 37036 |
| 20 | 16550 | 20089 | 9762 | 10608 | 14077 | 22171 | 26974 | 48585 | 13949 | 27586 | 37391 | 19179 |
| 21 | 8641 | 10071 | 15293 | 8006 | 8654 | 11971 | 18871 | 22424 | 42543 | 11600 | 24431 | 31974 |
| 22 | 4570 | 6404 | 7446 | 13309 | 6507 | 7296 | 10024 | 15752 | 19425 | 48332 | 10300 | 21626 |
| 23 | 22539 | 4021 | 4855 | 6325 | 11316 | 5435 | 5929 | 7973 | 13537 | 18218 | 43025 | 8515 |
| 24 | 16632 | 19656 | 2975 | 3602 | 5249 | 9794 | 4456 | 4495 | 6711 | 12943 | 18181 | 38202 |
| 25 | 25698 | 13570 | 17096 | 2511 | 2885 | 4435 | 8388 | 3313 | 3687 | 5503 | 11487 | 14265 |

Number of Animals 2312753 2145031 2066806 1932953 1823934 1768570 1743757 1597515 1554184 1532589 1556169 1335783

Table 5. Numbers at age of the northwest Atlantic harp seal estimated by cohort analysis from catch data (Table 3) assuming $M=0.144$.

| Age | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 444307 | 343878 | 373190 | 458617 | 412971 | 445669 | 420976 | 380759 | 350972 | 313938 | 358976 | 376386 |
| 1 | 134689 | 140497 | 129826 | 120212 | 144853 | 217369 | 174225 | 167317 | 138207 | 200129 | 183062 | 208804 |
| 2 | 150394 | 117814 | 118357 | 104921 | 96986 | 123598 | 174127 | 148044 | 142380 | 122381 | 176726 | 154267 |
| 3 | 139722 | 130017 | 101632 | 95797 | 88799 | 82266 | 107454 | 148249 | 129748 | 125823 | 106481 | 152368 |
| 4 | 110721 | 118656 | 111382 | 85786 | 83626 | 76333 | 70485 | 83313 | 129829 | 113623 | 109894 | 93003 |
| 5 | 100392 | 91774 | 99819 | 94267 | 74569 | 72894 | 65736 | 60394 | 82161 | 114355 | 99410 | 96399 |
| 6 | 76516 | 84509 | 74434 | 83510 | 80723 | 64946 | 62238 | 56398 | 52706 | 71793 | 98854 | 86862 |
| 7 | 77792 | 60997 | 68462 | 60666 | 70016 | 70409 | 55942 | 53939 | 49600 | 46044 | 62884 | 85233 |
| 8 | 47790 | 65522 | 51477 | 56011 | 50029 | 60171 | 60482 | 48237 | 47555 | 43482 | 39856 | 55179 |
| 9 | 42732 | 39412 | 56737 | 42584 | 47086 | 42279 | 50870 | 52419 | 42528 | 41721 | 37493 | 34429 |
| 10 | 34860 | 35116 | 34144 | 48718 | 35892 | 40403 | 35528 | 43458 | 46038 | 37323 | 36050 | 32242 |
| 11 | 36749 | 26617 | 29893 | 28644 | 41741 | 30365 | 34320 | 29851 | 37985 | 40283 | 32322 | 31265 |
| 12 | 27010 | 30337 | 23130 | 24290 | 23403 | 36006 | 25520 | 29408 | 26111 | 33333 | 34628 | 28148 |
| 13 | 32513 | 21697 | 25096 | 18979 | 20190 | 19907 | 31038 | 21512 | 25667 | 22738 | 28457 | 30055 |
| 14 | 30145 | 27099 | 18736 | 20844 | 15692 | 17093 | 16539 | 26584 | 18732 | 22380 | 19132 | 24612 |
| 15 | 48465 | 25019 | 22712 | 14993 | 16849 | 12836 | 13992 | 13960 | 23458 | 16378 | 18964 | 16424 |
| 16 | 36942 | 40599 | 20987 | 18805 | 11668 | 14011 | 10138 | 11341 | 12140 | 20543 | 13606 | 16284 |
| 17 | 14999 | 30776 | 35475 | 17233 | 15561 | 9431 | 11566 | 8182 | 9838 | 10545 | 17471 | 11492 |
| 18 | 36970 | 10656 | 26977 | 30610 | 13522 | 13032 | 7236 | 9520 | 7032 | 8529 | 8764 | 15028 |
| 19 | 15408 | 28935 | 8709 | 22541 | 25809 | 10896 | 10646 | 5720 | 8157 | 6063 | 7022 | 7398 |
| 20 | 31873 | 11546 | 24909 | 6793 | 18598 | 22014 | 8738 | 8835 | 4867 | 7129 | 4913 | 5909 |
| 21 | 15879 | 28305 | 9981 | 21192 | 4981 | 15889 | 18767 | 7158 | 7579 | 4104 | 5826 | 4095 |
| 22 | 27488 | 12100 | 24895 | 8169 | 18263 | 3986 | 13535 | 16878 | 6195 | 6638 | 3381 | 4932 |
| 23 | 18479 | 23455 | 10530 | 21705 | 6825 | 15649 | 3034 | 11782 | 14343 | 5439 | 5271 | 2745 |
| 24 | 6920 | 15421 | 20873 | 8747 | 18845 | 5732 | 13499 | 2532 | 10296 | 12734 | 4663 | 4513 |
| 25 | 33551 | 4648 | 13677 | 18272 | 7427 | 16562 | 4862 | 11861 | 2137 | 9136 | 11220 | 3954 |

Number 1+ 1328999 1221524 1062850 1074289 1031953 1094077 1080517 1076892 1075289 1142646 1166351 1205640
Animals

Table 6. Projected populations of male and female northwest Atlantic harp seals, and age specific sex ratios starting from 1961. The sex ratio of the large vessel catch was determined empirically, while the sex ratio of the landmen catch was considered to be that of the population. A maturity ogive was applied to calculate the breeding stock size.

| Age | 1962 | | 1963 | | 1964 | | 1965 | | 1966 | | 1967 | |
|-----|--------|-------------|-------|-------------|--------|-------------|-------|-------------|-------|-------------|-------|-------------|
| | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F |
| 1 | 107566 | 107566 | 87866 | 87866 | 67345 | 67345 | 70249 | 70249 | 64913 | 64913 | 60106 | 60106 |
| 2 | 93245 | 93245 | 84090 | 84090 | 75864 | 75864 | 59040 | 59040 | 59649 | 59649 | 53005 | 53005 |
| 3 | 71465 | 71465 | 68484 | 68484 | 71746 | 71746 | 65837 | 65826 | 51174 | 51156 | 48815 | 48792 |
| 4 | 55941 | 55956 | 59542 | 59660 | 57990 | 58204 | 61260 | 61415 | 56654 | 56813 | 43342 | 43554 |
| 5 | 55729 | 55765 | 50.02 | 45695 | 51.154 | 51482 | 50.16 | 48383 | 49042 | 50.34 | 51829 | 52381 |
| 6 | 34253 | 34273 | 50.01 | 46552 | 46811 | 50.14 | 39055 | 43162 | 43901 | 50.42 | 39434 | 41037 |
| 7 | 31154 | 31185 | 50.02 | 29235 | 29384 | 50.13 | 39578 | 31089 | 32562 | 51.16 | 34849 | 36819 |
| 8 | 26045 | 26090 | 50.04 | 26444 | 26617 | 50.16 | 24131 | 33258 | 34557 | 50.96 | 26205 | 28032 |
| 9 | 27458 | 27490 | 50.03 | 21821 | 22047 | 50.26 | 21564 | 19768 | 21149 | 51.69 | 28845 | 30167 |
| 10 | 20400 | 20428 | 50.04 | 23125 | 23352 | 50.24 | 17590 | 17573 | 19086 | 52.06 | 17109 | 18515 |
| 11 | 23260 | 23321 | 50.07 | 17587 | 17682 | 50.14 | 18479 | 13174 | 15090 | 53.39 | 14830 | 16632 |
| 12 | 22148 | 22183 | 50.04 | 20079 | 20227 | 50.18 | 13602 | 15135 | 16625 | 52.35 | 11444 | 13239 |
| 13 | 33214 | 33226 | 50.01 | 18793 | 18976 | 50.24 | 16379 | 10806 | 12170 | 52.97 | 12332 | 14297 |
| 14 | 25747 | 25761 | 50.01 | 29103 | 29182 | 50.07 | 15131 | 13568 | 14635 | 51.89 | 9376 | 10583 |
| 15 | 12019 | 12041 | 50.04 | 22254 | 22371 | 50.13 | 24291 | 12470 | 13631 | 52.22 | 11245 | 12649 |
| 16 | 25071 | 25074 | 50.00 | 9657 | 9816 | 50.41 | 18541 | 20170 | 21502 | 51.60 | 10368 | 11763 |
| 17 | 11583 | 11601 | 50.02 | 21998 | 22048 | 50.05 | 7036 | 15309 | 16386 | 51.70 | 17583 | 18950 |
| 18 | 21264 | 21279 | 50.02 | 9478 | 9633 | 50.41 | 18526 | 4737 | 6109 | 56.33 | 13406 | 14455 |
| 19 | 11107 | 11114 | 50.02 | 18551 | 18622 | 50.10 | 7740 | 14202 | 15686 | 52.48 | 3744 | 5241 |
| 20 | 18711 | 18722 | 50.01 | 9606 | 9659 | 50.14 | 15978 | 5664 | 6575 | 53.72 | 12143 | 13738 |
| 21 | 12238 | 12245 | 50.00 | 16010 | 16124 | 50.18 | 7973 | 14197 | 14449 | 50.44 | 4870 | 5772 |
| 22 | 5176 | 5176 | 50.00 | 10835 | 10853 | 50.04 | 13767 | 5931 | 6695 | 53.02 | 12334 | 12741 |
| 23 | 21523 | 21531 | 50.01 | 4287 | 4334 | 50.27 | 9271 | 11677 | 12217 | 51.13 | 5131 | 5904 |
| 24 | 8100 | 8103 | 50.01 | 19177 | 19184 | 50.01 | 3502 | 7666 | 8073 | 51.29 | 10396 | 10876 |
| 25 | 5771 | 5771 | 50.00 | 7143 | 7158 | 50.05 | 16850 | 2249 | 2778 | 55.26 | 6805 | 7168 |

| | | | | | | | | | | | |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Fraction males in catch | 0.550 | 0.550 | 0.65 | 0.65 | 0.65 | 0.65 | 0.65 | 0.75 | 0.75 | 0.75 | 0.75 |
| Fraction females | 0.500 | 0.501 | 0.505 | 0.505 | 0.509 | 0.511 | 0.517 | 0.517 | 0.517 | 0.517 | 0.517 |
| Fraction females in breeding population | 0.500 | 0.502 | 0.508 | 0.508 | 0.516 | 0.519 | 0.528 | 0.528 | 0.528 | 0.528 | 0.528 |
| Breeding population size | 424561 | 404631 | 384887 | 384887 | 356267 | 350906 | 338231 | 338231 | 338231 | 338231 | 338231 |

Table 6. Projected populations of male and female northwest Atlantic harp seals, and age specific sex ratios starting from 1961. The sex ratio of the large vessel catch was determined empirically, while the sex ratio of the landsmen catch was considered to be that of the population. A maturity ogive was applied to calculate the breeding stock size.

| Age | 1968 | | 1969 | | 1970 | | 1971 | | 1972 | | 1973 | |
|-----|-------|-------------|--------|-------------|-------|-------------|-------|-------------|-------|-------------|--------|-------------|
| | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F |
| 1 | 72427 | 72427 | 108685 | 108685 | 87113 | 87113 | 83659 | 83659 | 69104 | 69104 | 100065 | 100065 |
| 2 | 48961 | 48947 | 62195 | 62189 | 86146 | 86097 | 74467 | 74453 | 71782 | 71766 | 61388 | 61369 |
| 3 | 45115 | 45081 | 41856 | 41826 | 54241 | 54217 | 75554 | 75554 | 65503 | 65458 | 63512 | 63453 |
| 4 | 42689 | 42716 | 38904 | 39078 | 35964 | 36097 | 47232 | 47291 | 50.03 | 66365 | 58045 | 57983 |
| 5 | 37710 | 38025 | 37241 | 37480 | 33530 | 33932 | 30882 | 31225 | 50.28 | 41646 | 58486 | 58512 |
| 6 | 41074 | 42066 | 32786 | 33422 | 31703 | 32466 | 29407 | 30.51 | 27004 | 27429 | 36800 | 36924 |
| 7 | 36187 | 38221 | 35747 | 37044 | 28074 | 29208 | 27426 | 28433 | 50.90 | 25353 | 23704 | 24104 |
| 8 | 26253 | 29459 | 30835 | 33574 | 30442 | 32416 | 24019 | 25621 | 51.61 | 24171 | 22360 | 22992 |
| 9 | 23353 | 26994 | 21837 | 25840 | 25606 | 29348 | 26296 | 28432 | 51.95 | 21159 | 21296 | 22256 |
| 10 | 17761 | 20814 | 19742 | 23794 | 17981 | 22599 | 21707 | 25632 | 54.15 | 23025 | 18723 | 20137 |
| 11 | 20931 | 22946 | 14717 | 18273 | 16476 | 20825 | 14976 | 19610 | 56.70 | 18916 | 20370 | 22341 |
| 12 | 11272 | 13663 | 54.79 | 17831 | 12087 | 15944 | 14003 | 18214 | 56.54 | 13068 | 16580 | 19931 |
| 13 | 9546 | 12238 | 56.18 | 9417 | 15182 | 17826 | 10012 | 13897 | 58.13 | 12164 | 15231 | 17722 |
| 14 | 7402 | 9783 | 56.93 | 7882 | 10761 | 10449 | 12864 | 15615 | 54.83 | 8680 | 10713 | 14235 |
| 15 | 7864 | 10499 | 57.17 | 8541 | 6137 | 9373 | 6263 | 9110 | 59.26 | 11339 | 7465 | 10656 |
| 16 | 5412 | 7536 | 58.20 | 6317 | 4245 | 7367 | 4769 | 8067 | 62.85 | 5411 | 9926 | 12199 |
| 17 | 7378 | 9362 | 55.93 | 4154 | 4988 | 8011 | 3265 | 6339 | 66.00 | 4092 | 4723 | 7094 |
| 18 | 6222 | 8509 | 57.76 | 6009 | 8195 | 5632 | 3974 | 6924 | 63.54 | 2755 | 3485 | 6203 |
| 19 | 12444 | 14434 | 53.70 | 4720 | 4653 | 7144 | 2093 | 4849 | 69.85 | 3334 | 2266 | 4795 |
| 20 | 8781 | 10789 | 55.13 | 10383 | 3488 | 6464 | 3741 | 6203 | 62.38 | 1722 | 2887 | 5355 |
| 21 | 1741 | 3644 | 67.67 | 7331 | 8615 | 11224 | 2731 | 5589 | 67.18 | 3148 | 1248 | 3540 |
| 22 | 8651 | 10536 | 54.91 | 1229 | 6074 | 8395 | 7457 | 9923 | 57.09 | 2317 | 2642 | 4709 |
| 23 | 3143 | 4315 | 57.86 | 7261 | 708 | 2760 | 5230 | 7422 | 58.66 | 6468 | 8815 | 1954 |
| 24 | 9292 | 9932 | 51.66 | 2546 | 6129 | 8236 | 531 | 2412 | 81.98 | 4521 | 5728 | 7825 |
| 25 | 3473 | 4447 | 56.15 | 8108 | 2081 | 3357 | 5354 | 7302 | 57.70 | 391 | 3964 | 5819 |

Fraction males in catch 0.75 0.86 0.86 0.86 0.86 0.86

Fraction females 0.520 0.521 0.524 0.524 0.524 0.518

Fraction females in breeding population 0.534 0.541 0.549 0.549 0.545 0.538

Breeding population size 321977 315332 302092 296103 294071 299838

Table 6. Projected populations of male and female northwest Atlantic harp seals, and age specific sex ratios starting from 1961. The sex ratio of the large vessel catch was determined empirically, while the sex ratio of the landsmen catch was considered to be that of the population. A maturity ogive was applied to calculate the breeding stock size.

| Age | 1974 | | 1975 | | 1976 | | 1977 | |
|---|--------|-------------|--------|-------------|--------|-------------|--------|-------------|
| | Males | Females & F | Males | Females & F | Males | Females & F | Males | Females & F |
| 1 | 91531 | 50.00 | 104402 | 50.00 | 73598 | 50.00 | 79888 | 50.00 |
| 2 | 88545 | 50.00 | 77800 | 49.98 | 88916 | 49.96 | 61790 | 49.93 |
| 3 | 53788 | 53656 | 76733 | 76697 | 66909 | 66743 | 77242 | 77002 |
| 4 | 55636 | 55739 | 46964 | 47090 | 66984 | 66982 | 58097 | 49.93 |
| 5 | 50844 | 51096 | 48811 | 49142 | 40965 | 41165 | 58725 | 58724 |
| 6 | 50507 | 51322 | 44389 | 44987 | 42565 | 42978 | 35969 | 36176 |
| 7 | 32220 | 32616 | 43466 | 44838 | 38682 | 39393 | 37512 | 37898 |
| 8 | 20512 | 21158 | 28255 | 28796 | 37919 | 39231 | 34100 | 34785 |
| 9 | 19217 | 20199 | 17704 | 18502 | 24628 | 25214 | 33440 | 34653 |
| 10 | 18425 | 19489 | 16424 | 17703 | 15390 | 16253 | 21779 | 22345 |
| 11 | 16219 | 17666 | 15938 | 17117 | 14240 | 15459 | 13449 | 14233 |
| 12 | 17536 | 19515 | 14087 | 15553 | 13846 | 15045 | 12507 | 13612 |
| 13 | 14209 | 17335 | 15151 | 17185 | 12256 | 13648 | 12212 | 13299 |
| 14 | 9610 | 13212 | 12215 | 15261 | 13179 | 15095 | 10808 | 12083 |
| 15 | 9100 | 12352 | 8190 | 11617 | 55.54 | 53.39 | 11638 | 13365 |
| 16 | 6166 | 9192 | 7734 | 10860 | 58.41 | 55.81 | 9287 | 11792 |
| 17 | 8431 | 10620 | 5135 | 8011 | 60.94 | 58.78 | 6265 | 9084 |
| 18 | 3892 | 6134 | 7192 | 9323 | 56.45 | 44.36 | 7073 | 8466 |
| 19 | 2815 | 5356 | 3212 | 5370 | 62.57 | 62.48 | 8200 | 58.94 |
| 20 | 1746 | 4155 | 2291 | 4694 | 67.20 | 2774 | 4742 | 61.63 |
| 21 | 2273 | 4651 | 1377 | 3640 | 72.56 | 1972 | 4148 | 56.83 |
| 22 | 952 | 3093 | 1869 | 4080 | 68.58 | 1175 | 3222 | 63.33 |
| 23 | 2053 | 3982 | 682 | 2695 | 79.80 | 1601 | 1743 | 67.91 |
| 24 | 1612 | 3828 | 1710 | 3511 | 67.25 | 569 | 1027 | 2862 |
| 25 | 5012 | 6966 | 1286 | 3391 | 72.51 | 1466 | 1413 | 3211 |
| | | | | | | | 497 | 2123 |
| | | | | | | | | 81.04 |
| Fraction males in catch | 0.86 | | 0.86 | | 0.86 | | 0.86 | |
| Fraction females | 0.517 | | 0.516 | | 0.514 | | 0.512 | |
| Fraction females in breeding population | 0.536 | | 0.534 | | 0.529 | | 0.524 | |
| Breeding population size | 302165 | | 305609 | | 311502 | | 319914 | |




Figure 1. The fraction of males in the Norwegian catch of moulted northwest Atlantic harp seals from 1969 to 1974 (from Benjaminson and Øritsland MS 1975)




Figure 2. The percentage landmen hunt of one year and older harp seals from 1952 to 1975.




Figure 3. A comparison of three different methods of estimating pup production.




Figure 4. A shifting maturity ogive in relation to population size. The ogive is constrained in the model so it cannot shift further to the left than the 1.2×10^{-6} ogive although the population may continue to decline.




Figure 5. The decline in reproductive successful northwest Atlantic female harp seals in relation to population size.

```

      VVARP(0)
V HARP
[1] 'STARTING POP ONE AND OLDER'
[2] POP1←POP-0
[3] 'NUMBER OF YEARS'
[4] Y←0
[5] 'ENTER FIRST YEAR'
[6] DAY←DATE-0
[7] 'LARGE VESSEL CATCH'
[8] LVC←0
[9] 'POP CATCH'
[10] PC←0
[11] 'MAGDALLEN QUOTA ON PUPS'
[12] MPQ←0
[13] 'MAGDALLEN QUOTA ON ONE PLUS'
[14] MAQ←0
[15] 'NUMBER OF RUNS'
[16] Z←0
[17] STAR←0
[18] BPOP←YEAR←TPOP←TSRX←BSRX←TCA←TCP←10
[19] →START
[20] TAKEOFF←(QM<23.50)/BACK
[21] MAT← 0 0 0.16 0.4 0.66 0.89 0.97 ,(18p1)
[22] →AA
[23] FIX1:QM←0
[24] →LEFT
[25] FIX2:QM←90
[26] →LEFT
[27] AAAA:BRRED←0
[28] →ABC
[29] BBBB:TOTALP←0
[30] →BCD
[31] START:STAR←STAR+1
[32] POP←POP1
[33] DAY←DAY
[34] CR← 0.5 0.4993 0.4992 0.4993 0.4999 0.5014 0.5026 0.505 0.5089 0.5064 0.5142 0.5211 0.5213
    0.5278 0.5345
[35] CR←CR, 0.5594 0.5918 0.5894 0.8183 0.5683 0.8333 0.8791 0.7359 0.8945 0.8104
[36] PPOP←POP×CR
[37] MPOP←POP-PPOP
[38] →AROUND
[39] GO1:PRRG←1
[40] →SKIP
[41] XK←0
[42] AROUND:XX←YY+0
[43] GO2:XY←XX+1
[44] DATE←DATE+1
[45] CORR← 0.528 0.528 0.591 0.845 0.693 0.731 0.764 0.79 0.811 0.826 0.838 0.846 0.851 0.853 ,(11p
    0.854)
[46] PLV← 0.3071 0.14 0.077 0.0602 0.049 0.042 0.0364 0.0328 0.0294 0.0272 0.0248 0.227 0.021
    0.0196 0.0182
[47] PLV←PLV, 0.0174 0.0154 0.0134 0.0126 0.0101 0.0078 0.0067 0.005 0.0042 0.0008
[48] PLC← 0.1187 0.107 0.0978 0.0877 0.0781 0.0701 0.0618 0.0544 0.048 0.0424 0.0369 0.0323 0.0277
    0.024 0.0207
[49] PLC←PLC, 0.01077 0.0151 0.0129 0.0109 0.0092 0.0078 0.0063 0.0052 0.0041 0.0031
[50] PGC← 0.3059 0.1851 0.1253 0.0839 0.0575 0.0425 0.031 0.0241 0.0184 0.0149 0.0128 0.0108 0.0097
    0.0086 0.0078
[51] PGC←PGC, 0.0074 0.0069 0.0067 0.0063 0.0061 0.0059 0.0057 0.0056 0.0055 0.0054
[52] PNC← 0.1312 0.1602 0.2123 0.1158 0.083 0.0617 0.0483 0.0367 0.0289 0.0232 0.0183 0.0154 0.0125
    0.0104 0.0085
[53] PNC←PNC, 0.0071 0.0058 0.005 0.0042 0.0034 0.0031 0.0027 0.0023 0.0019 0
[54] MLVC←LVC×PLV×CORR
[55] PLVC←(LVC×PLV)-MLVC
[56] →(MAQ>0)/YY
[57] LC←(5048×((+/(1271000)+1000))-8))+13026
[58] →BBB
[59] YY:LC←((5051×((+/(1271000)+1000))-8))+12451+MAQ
[60] BBB←PF←PPOP+(FPOP+MPOP)
[61] PLC←PF×PLC×LC
[62] NLC←(PLC×LC)-PLC
[63] GC←(1040×((+/(1271000)+1000))-8))+3784
[64] FGC←PF×PGC×GC
[65] NGC←(PGC×GC)-FGC
[66] NC←(729×((+/(1271000)+1000))-8))+1294
[67] FNC←PF×PNC×NC
[68] NNC←(PNC×NC)-FNC
[69] POPU←/(~24+POP)
[70] PRRG←(1.048-(0.097458×1E~6×POPU))
[71] →(PRRG>1)/GO1
[72] SKIP:AGR←125
[73] M←(15.5223×AGR)+(~2.245E~5×(+/POP))+(~16.01743)
[74] MAT←10
[75] GOTO:YY←YY+1
[76] QM←1+M
[77] →(YY=4)/TAKEOFF
[78] BACK:→(QM<0)/FIX1
[79] →(QM>90)/FIX2

```

Figure 6. Program listing of APL simulation of north-west Atlantic harp seal.

```

[80] LETT:QW=10(QW+57.296)
[81] NAT=NAT,QW
[82] N=1+N
[83] →(YY<25)/GOTO
[84] AA:PUP→BREEDPOP+(+/(FPOP=PREG=NAT))*1.06
[85] YY=0
[86] FPOP→FPOP-(FLVC+PLC+PGC+PNC)
[87] NPOP→NPOP-(MLVC+NLC+NGC+NMC)
[88] TAC→+/(FLVC+PLC+PGC+PNC+MLVC+NLC+NGC+NMC)
[89] →(NPQ=0)/XX
[90] LPC=(14442*((+/(1271000)+1000))-6)+36949
[91] →AAA
[92] XX:LPC+((13998*((+/(1271000)+1000))-6))+24177)+NPQ
[93] AAA:GPC+(1040*((+/(1271000)+1000))-6)+3784
[94] PUP→PUP-(PC+LPC+GPC)
[95] TPC→PC+LPC+GPC
[96] NMORT=(0.0302*((+/(1271000)+1000))-6)+0.114
[97] FPOP→FPOP*(+(-NMORT))
[98] NPOP→NPOP*(+(-NMORT))
[99] PUP→PUP*(+(-NMORT))
[100] ONE→PUP+2
[101] FPOP→ONE,24,FPOP
[102] NPOP→ONE,24,NPOP
[103] POP→FPOP+NPOP
[104] BREED→[+/(BREEDPOP)]
[105] →(BREED<0)/AAAA
[106] ABC:BPOP→BPOP,BREED
[107] TOTALP→[+/(POP)]
[108] →(TOTALP<0)/BBBB
[109] BCD:TPOP→TPOP,TOTALP
[110] SEXRT→[(((+/(FPOP)+(+/(POP)))×100)]
[111] SEXRB→[(((+/(FPOP=NAT))+(+/(POP=NAT)))×100)]
[112] TCA→TCA,TAC
[113] TCP→TCP,TPC
[114] TSEX→TSEX,SEXRT
[115] BSEX→BSEX,SEXRB
[116] YEAR→YEAR,DATE
[117] →(X<X)/GO2
[118] →(Z=1)/GO3
[119] DATA→[Q(7,X)P(YEAR,BPOP,TPOP,TSEX,BSEX,TCA,TCP)]
[120] DATA
[121] →0
[122] GO3:→(Z>STAR)/START
[123] BPOP→[Q((Z,X)P(BPOP)]
[124] TPOP→[Q((Z,X)P(TPOP)]
[125] TCA→[Q((Z,X)P(TCA)]
[126] TCP→[Q((Z,X)P(TCP)]
[127] X=Z
[128] NBPOP→(+/[2] BPOP)+X
[129] SDBPOP→(((+/[2])(BPOP)*2))-(Z*(NBPOP*2)))+(Z-1)*0.5
[130] NATBPOP→BPOP,NBPOP
[131] NATBPOP→[ (NATBPOP,SDBPOP) ]
[132] NTPOP→(+/[2] TPOP)+X
[133] SDTPOP→(((+/[2])(TPOP)*2))-(Z*(NTPOP*2)))+(Z-1)*0.5
[134] NATTPOP→TPOP,NTPOP
[135] NATTPOP→[ (NATTPOP,SDTPOP) ]
[136] NTCA→(+/[2] TCA)+X
[137] SDTCA→(((+/[2])(TCA)*2))-(Z*(NTCA*2)))+(Z+1)*0.5
[138] NATTCA→TCA,NTCA
[139] NATTCA→[ (NATTCA,SDTCA) ]
[140] NTCP→(+/[2] TCP)+X
[141] SDTCP→(((+/[2])(TCP)*2))-(Z*(NTCP*2)))+(Z+1)*0.5
[142] NATTCP→TCP,NTCP
[143] NATTCP→[ (NATTCP,SDTCP) ]
[144] ' TOTAL POPULATION '
[145] NATPOP
[146] ''
[147] ''
[148] ' BREEDING POPULATION '
[149] NATBPOP
[150] ''
[151] ''
[152] ' CATCH OF PUPS '
[153] NATTCP
[154] ''
[155] ''
[156] ' CATCH OF ADULTS '
[157] NATTCA

```

Figure 6. Program listing of APL simulation of north-west Atlantic harp seal.




Figure 8. Catch distribution from the different fisheries for the northwest Atlantic harp seal.




Figure 9. Two simulated relationships showing the effects of density dependent age of maturity and pregnancy rates on the northwest Atlantic harp seal.




Figure 10. Simulated recruitment curve, and pup production from cohort analysis of the north-west Atlantic harp seal. Bars represent 2 (SD).




Figure 11. Shaefer type curve illustrating the MSY population level of ~ 1.6 million for catch of pups and 1+ northwest Atlantic harp seal in their present proportions. Since lower catches of pups allows for larger population sizes a constant hunting mortality by large vessel will allow for a greater catch of 1+ seals. Bars represent $2(SD)$.




Figure 12. Projected breeding stock of the northwest Atlantic harp seal in relation to varying management strategies. Confidence limits indicate we can put little reliance in projections further than 5 years.